

**GPR18 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51240****Specification**

---

**GPR18 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q14330</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	38 KDa
Antigen Region	121 - 180

**GPR18 Antibody - Additional Information****Gene ID** 2841**Other Names**

N-arachidonyl glycine receptor, NAGly receptor, G-protein coupled receptor 18, GPR18, GPCRW

**Target/Specificity**

KLH conjugated synthetic peptide derived from human GPR18

**Dilution**

WB~~ 1:1000

**Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**GPR18 Antibody - Protein Information****Name** GPR18**Synonyms** GPCRW**Function**

Receptor for endocannabinoid N-arachidonyl glycine (NAGly) (PubMed:<a href="http://www.uniprot.org/citations/16844083" target="\_blank">16844083</a>, PubMed:<a href="http://www.uniprot.org/citations/24762058" target="\_blank">24762058</a>, PubMed:<a href="http://www.uniprot.org/citations/27572937" target="\_blank">27572937</a>). However, conflicting results about the role of NAGly as an agonist are reported (PubMed:<a href="http://www.uniprot.org/citations/27018161" target="\_blank">27018161</a>). Can also be activated by plant-derived and synthetic cannabinoid agonists (PubMed:<a href="http://www.uniprot.org/citations/24762058" target="\_blank">24762058</a>). The activity

of this receptor is mediated by G proteins which inhibit adenylyl cyclase (PubMed:<a href="http://www.uniprot.org/citations/16844083" target="\_blank">16844083</a>). May contribute to regulation of the immune system. Is required for normal homeostasis of CD8+ subsets of intraepithelial lymphocytes (IELs) (CD8alphaalpha and CD8alphabeta IELs)in small intestine by supporting preferential migration of CD8alphaalpha T-cells to intraepithelial compartment over lamina propria compartment, and by mediating their reconstitution into small intestine after bone marrow transplant (By similarity). Plays a role in hypotensive responses, mediating reduction in intraocular and blood pressure (By similarity). Mediates NAGly-induced process of reorganization of actin filaments and induction of acrosomal exocytosis (PubMed:<a href="http://www.uniprot.org/citations/27572937" target="\_blank">27572937</a>).

#### Cellular Location

Cell membrane; Multi-pass membrane protein. Cytoplasmic vesicle membrane

#### Tissue Location


Expressed in midpiece of spermatozoon (at protein level) (PubMed:27572937). Most abundant in testis and spleen (PubMed:16844083). Highly expressed in CD4 and CD8-positive T-cells as well as CD19-positive B-cells (PubMed:16844083)

### GPR18 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### GPR18 Antibody - Images



Anti-GPR18 Antibody at 1:1000 dilution + Jurkat whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 38 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

**GPR18 Antibody - Background**

Receptor for N-arachidonyl glycine. The activity of this receptor is mediated by G proteins which inhibit adenylyl cyclase. May contribute to regulation of the immune system.

**GPR18 Antibody - References**

Gantz I.,et al.Genomics 42:462-466(1997).  
Kohno M.,et al.Biochem. Biophys. Res. Commun. 347:827-832(2006).  
Xu X.,et al.Submitted (MAY-2000) to the EMBL/GenBank/DDBJ databases.  
Kalnine N.,et al.Submitted (AUG-2003) to the EMBL/GenBank/DDBJ databases.  
Dunham A.,et al.Nature 428:522-528(2004).