

## **HSPE1 Antibody**

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51272

### **Specification**

## **HSPE1 Antibody - Product Information**

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW
Antigen Region

WB
P61604
Human, Mouse, Rat
Rabbit
Polyclonal
10 KDa
41 - 100

# **HSPE1** Antibody - Additional Information

**Gene ID 3336** 

#### **Other Names**

10 kDa heat shock protein, mitochondrial, Hsp10, 10 kDa chaperonin, Chaperonin 10, CPN10, Early-pregnancy factor, EPF, HSPE1

# **Target/Specificity**

KLH conjugated synthetic peptide derived from human HSPE1

#### **Dilution**

WB~~ 1:1000

#### **Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

# **Storage**

Store at -20 °C. Stable for 12 months from date of receipt

# **HSPE1 Antibody - Protein Information**

# Name HSPE1

#### **Function**

Co-chaperonin implicated in mitochondrial protein import and macromolecular assembly. Together with Hsp60, facilitates the correct folding of imported proteins. May also prevent misfolding and promote the refolding and proper assembly of unfolded polypeptides generated under stress conditions in the mitochondrial matrix (PubMed:<a

 $href="http://www.uniprot.org/citations/11422376" target="\_blank">11422376</a>, PubMed: <a href="http://www.uniprot.org/citations/1346131" target="\_blank">1346131</a>, PubMed: <a href="http://www.uniprot.org/citations/7912672" target="_blank">7912672</a>). The functional units of these chaperonins consist of heptameric rings of the large subunit Hsp60, which function as a back-to-back double ring. In a cyclic reaction, Hsp60 ring complexes bind one unfolded$ 



substrate protein per ring, followed by the binding of ATP and association with 2 heptameric rings of the co-chaperonin Hsp10. This leads to sequestration of the substrate protein in the inner cavity of Hsp60 where, for a certain period of time, it can fold undisturbed by other cell components. Synchronous hydrolysis of ATP in all Hsp60 subunits results in the dissociation of the chaperonin rings and the release of ADP and the folded substrate protein (Probable).


**Cellular Location**Mitochondrion matrix.

### **HSPE1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### **HSPE1 Antibody - Images**




All lanes : Anti-HSPE1 Antibody at 1:1000 dilution Lane 1: Jurkat whole cell lysates Lane 2: Hela whole cell lysates Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 11 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

# **HSPE1 Antibody - Background**

Eukaryotic CPN10 homolog which is essential for mitochondrial protein biogenesis, together with CPN60. Binds to CPN60 in the presence of Mg-ATP and suppresses the ATPase activity of the latter.

## **HSPE1 Antibody - References**

Monzini N., et al. Biochim. Biophys. Acta 1218:478-480(1994).





Chen J.J., et al. Biochim. Biophys. Acta 1219:189-190(1994). Hansen J.J., et al. Hum. Genet. 112:71-77(2003). Ota T., et al. Nat. Genet. 36:40-45(2004). Ebert L., et al. Submitted (MAY-2004) to the EMBL/GenBank/DDBJ databases.