

## MDA5 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51277

## Specification

# MDA5 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW

WB, ICC, IHC-P, E <u>Q9BYX4</u> Human, Rat Rabbit Polyclonal 17 KDa

## **MDA5 Antibody - Additional Information**

Gene ID 64135

#### **Other Names**

Interferon-induced helicase C domain-containing protein 1, Clinically amyopathic dermatomyositis autoantigen 140 kDa, CADM-140 autoantigen, Helicase with 2 CARD domains, Helicard, Interferon-induced with helicase C domain protein 1, Melanoma differentiation-associated protein 5, MDA-5, Murabutide down-regulated protein, RIG-I-like receptor 2, RLR-2, RNA helicase-DEAD box protein 116, IFIH1, MDA5, RH116

Dilution WB~~1:1000 ICC~~N/A IHC-P~~N/A E~~N/A

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

# MDA5 Antibody - Protein Information

Name IFIH1 (<u>HGNC:18873</u>)

#### Function

Innate immune receptor which acts as a cytoplasmic sensor of viral nucleic acids and plays a major role in sensing viral infection and in the activation of a cascade of antiviral responses including the induction of type I interferons and pro-inflammatory cytokines (PubMed:<a href="http://www.uniprot.org/citations/28594402" target="\_blank">28594402</a>, PubMed:<a href="http://www.uniprot.org/citations/28594402" target="\_blank">28594402</a>, PubMed:<a href="http://www.uniprot.org/citations/32169843" target="\_blank">32169843</a>, PubMed:<a href="http://www.uniprot.org/citations/32169843" target="\_blank">33727702</a>). Its ligands include mRNA lacking 2'-O-methylation at their 5' cap and long-dsRNA (>1 kb in length)



(PubMed:<a href="http://www.uniprot.org/citations/22160685" target=" blank">22160685</a>). Upon ligand binding it associates with mitochondria antiviral signaling protein (MAVS/IPS1) which activates the IKK-related kinases: TBK1 and IKBKE which phosphorylate interferon regulatory factors: IRF3 and IRF7 which in turn activate transcription of antiviral immunological genes, including interferons (IFNs); IFN-alpha and IFN-beta. Responsible for detecting the Picornaviridae family members such as encephalomyocarditis virus (EMCV), mengo encephalomyocarditis virus (ENMG), and rhinovirus (PubMed:<a href="http://www.uniprot.org/citations/28606988" target=" blank">28606988</a>). Detects coronavirus SARS-CoV-2 (PubMed:<a href="http://www.uniprot.org/citations/33440148" target=" blank">33440148</a>, PubMed:<a href="http://www.uniprot.org/citations/33514628" target="\_blank">33514628</a>). Can also detect other viruses such as dengue virus (DENV), west Nile virus (WNV), and reovirus. Also involved in antiviral signaling in response to viruses containing a dsDNA genome, such as vaccinia virus. Plays an important role in amplifying innate immune signaling through recognition of RNA metabolites that are produced during virus infection by ribonuclease L (RNase L). May play an important role in enhancing natural killer cell function and may be involved in growth inhibition and apoptosis in several tumor cell lines.

#### **Cellular Location**

Cytoplasm. Nucleus. Mitochondrion. Note=Upon viral RNA stimulation and ISGylation, translocates from cytosol to mitochondrion. May be found in the nucleus, during apoptosis

#### **Tissue Location**

Widely expressed, at a low level. Expression is detected at slightly highest levels in placenta, pancreas and spleen and at barely levels in detectable brain, testis and lung

## MDA5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

#### MDA5 Antibody - Images

## MDA5 Antibody - Background

Innate immune receptor which acts as a cytoplasmic sensor of viral nucleic acids and plays a major role in sensing viral infection and in the activation of a cascade of antiviral responses including the induction of type I interferons and proinflammatory cytokines. Its ligands include mRNA lacking 2'-O- methylation at their 5' cap and long-dsRNA (>1 kb in length). Upon ligand binding it associates with mitochondria antiviral signaling protein (MAVS/IPS1) which activates the IKK-related kinases: TBK1 and IKBKE which phosphorylate interferon regulatory factors: IRF3 and IRF7 which in turn activate transcription of antiviral immunological genes, including interferons (IFNs); IFN-alpha and IFN-beta. Responsible for detecting the Picornaviridae family members such as encephalomyocarditis virus (EMCV) and mengo encephalomyocarditis virus (ENMG). Can also detect other viruses such as dengue virus (DENV), west Nile virus (WNV), and reovirus. Also involved in antiviral signaling in response to viruses containing a dsDNA genome, such as vaccinia virus. Plays an important role in amplifying innate immune signaling through recognition of RNA metabolites that are produced during virus infection by ribonuclease L (RNase L). May play an important role in enhancing natural killer cell function and may be involved in growth inhibition and



apoptosis in several tumor cell lines.

## **MDA5 Antibody - References**

Kang D.-C., et al. Proc. Natl. Acad. Sci. U.S.A. 99:637-642(2002). Cocude C., et al.J. Gen. Virol. 84:3215-3225(2003). Ota T., et al.Nat. Genet. 36:40-45(2004). Kovacsovics M., et al. Curr. Biol. 12:838-843(2002). Kovacsovics M., et al. Curr. Biol. 12:1633-1633(2002).