

#### c-Maf Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51325

### **Specification**

### c-Maf Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, IP, E
O75444
Human, Mouse, Rat
Rabbit
Polyclonal
50 KDa

# c-Maf Antibody - Additional Information

# **Gene ID** 4094

#### **Other Names**

Transcription factor Maf, Proto-oncogene c-Maf, V-maf musculoaponeurotic fibrosarcoma oncogene homolog, MAF

### Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human c-Maf. The exact sequence is proprietary.

#### **Dilution**

WB~~1:1000 IP~~N/A E~~N/A

#### **Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

### **Storage**

Store at -20 °C. Stable for 12 months from date of receipt

## c-Maf Antibody - Protein Information

### **Name MAF**

#### **Function**

Acts as a transcriptional activator or repressor. Involved in embryonic lens fiber cell development. Recruits the transcriptional coactivators CREBBP and/or EP300 to crystallin promoters leading to up- regulation of crystallin gene during lens fiber cell differentiation. Activates the expression of IL4 in T helper 2 (Th2) cells. Increases T- cell susceptibility to apoptosis by interacting with MYB and decreasing BCL2 expression. Together with PAX6, transactivates strongly the glucagon gene promoter through the G1 element. Activates transcription of the CD13 proximal promoter in endothelial cells. Represses transcription of the CD13 promoter in early stages of myelopoiesis by



affecting the ETS1 and MYB cooperative interaction. Involved in the initial chondrocyte terminal differentiation and the disappearance of hypertrophic chondrocytes during endochondral bone development. Binds to the sequence 5'-[GT]G[GC]N[GT]NCTCAGNN-3' in the L7 promoter. Binds to the T-MARE (Maf response element) sites of lens-specific alpha- and beta-crystallin gene promoters. Binds element G1 on the glucagon promoter. Binds an AT-rich region adjacent to the TGC motif (atypical Maf response element) in the CD13 proximal promoter in endothelial cells (By similarity). When overexpressed, represses anti-oxidant response element (ARE)-mediated transcription. Involved either as an oncogene or as a tumor suppressor, depending on the cell context. Binds to the ARE sites of detoxifying enzyme gene promoters.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00978}.

**Tissue Location** 

Expressed in endothelial cells.

#### c-Maf Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

### c-Maf Antibody - Images

# c-Maf Antibody - Background

Acts as a transcriptional activator or repressor. Involved in embryonic lens fiber cell development. Recruits the transcriptional coactivators CREBBP and/or EP300 to crystallin promoters leading to up-regulation of crystallin gene during lens fiber cell differentiation. Activates the expression of IL4 in T helper 2 (Th2) cells. Increases T-cell susceptibility to apoptosis by interacting with MYB and decreasing BCL2 expression. Together with PAX6, transactivates strongly the glucagon gene promoter through the G1 element. Activates transcription of the CD13 proximal promoter in endothelial cells. Represses transcription of the CD13 promoter in early stages of myelopoiesis by affecting the ETS1 and MYB cooperative interaction. Involved in the initial chondrocyte terminal differentiation and the disappearance of hypertrophic chondrocytes during endochondral bone development. Binds to the sequence 5'-[GT]G[GC]N[GT]NCTCAGNN-3' in the L7 promoter. Binds to the T-MARE (Maf response element) sites of lens-specific alpha- and beta-crystallin gene promoters. Binds element G1 on the glucagon promoter. Binds an AT-rich region adjacent to the TGC motif (atypical Maf response element) in the CD13 proximal promoter in endothelial cells (By similarity). When overexpressed, represses anti-oxidant response element (ARE)- mediated transcription. Involved either as an oncogene or as a tumor suppressor, depending on the cell context. Binds to the ARE sites of detoxifying enzyme gene promoters.


# c-Maf Antibody - References

Chesi M., et al. Blood 91:4457-4463(1998).

Martin J., et al. Nature 432:988-994(2004).

Dhakshinamoorthy S., et al. Oncogene 21:5301-5312(2002).

Hurt E.M., et al. Cancer Cell 5:191-199(2004).





Watson J.E., et al. Oncogene 23:3487-3494(2004).