

Menin Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51340**Specification**

Menin Antibody - Product Information

Application	WB, IHC-P, E
Primary Accession	O00255
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	68 KDa

Menin Antibody - Additional Information**Gene ID** 4221**Other Names**

Menin, MEN1, SCG2

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Menin. The exact sequence is proprietary.

Dilution

WB~~1:1000

IHC-P~~N/A

E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

Menin Antibody - Protein Information**Name** MEN1**Synonyms** SCG2**Function**

Essential component of a MLL/SET1 histone methyltransferase (HMT) complex, a complex that specifically methylates 'Lys-4' of histone H3 (H3K4). Functions as a transcriptional regulator. Binds to the TERT promoter and represses telomerase expression. Plays a role in TGFB1-mediated inhibition of cell-proliferation, possibly regulating SMAD3 transcriptional activity. Represses JUND-mediated transcriptional activation on AP1 sites, as well as that mediated by NFkB subunit RELA. Positively regulates HOXC8 and HOXC6 gene expression. May be involved in normal

hematopoiesis through the activation of HOXA9 expression (By similarity). May be involved in DNA repair.

Cellular Location

Nucleus. Note=Concentrated in nuclear body-like structures. Relocates to the nuclear matrix upon gamma irradiation

Tissue Location

Ubiquitous.

Menin Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Menin Antibody - Images**Menin Antibody - Background**

Essential component of a MLL/SET1 histone methyltransferase (HMT) complex, a complex that specifically methylates 'Lys-4' of histone H3 (H3K4). Functions as a transcriptional regulator. Binds to the TERT promoter and represses telomerase expression. Plays a role in TGFB1-mediated inhibition of cell-proliferation, possibly regulating SMAD3 transcriptional activity. Represses JUND-mediated transcriptional activation on AP1 sites, as well as that mediated by NFkB subunit RELA. Positively regulates HOXC8 and HOXC6 gene expression. May be involved in normal hematopoiesis through the activation of HOXA9 expression (By similarity). May be involved in DNA repair.

Menin Antibody - References

Chandrasekharappa S.C.,et al.Science 276:404-407(1997).
Taylor T.D.,et al.Nature 440:497-500(2006).
Toledo R.A.,et al.Clin. Endocrinol. (Oxf.) 67:377-384(2007).
Agarwal S.K.,et al.Cell 96:143-152(1999).
Heppner C.,et al.Oncogene 20:4917-4925(2001).