

**MMP13 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51348****Specification**

---

**MMP13 Antibody - Product Information**

Application	WB, ICC, IHC-P, E
Primary Accession	<a href="#">P45452</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	48 KDa

**MMP13 Antibody - Additional Information****Gene ID** 4322**Other Names**

Collagenase 3, 3424-, Matrix metalloproteinase-13, MMP-13, MMP13

**Target/Specificity**

KLH-conjugated synthetic peptide encompassing a sequence within the N-term region of human MMP13. The exact sequence is proprietary.

**Dilution**

WB~~1:1000

ICC~~N/A

IHC-P~~N/A

E~~N/A

**Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**MMP13 Antibody - Protein Information****Name** MMP13**Function**

Plays a role in the degradation of extracellular matrix proteins including fibrillar collagen, fibronectin, TNC and ACAN. Cleaves triple helical collagens, including type I, type II and type III collagen, but has the highest activity with soluble type II collagen. Can also degrade collagen type IV, type XIV and type X. May also function by activating or degrading key regulatory proteins, such as TGFB1 and CCN2. Plays a role in wound healing, tissue remodeling, cartilage degradation, bone development, bone mineralization and ossification. Required for normal embryonic bone development and ossification. Plays a role in the healing of bone fractures via endochondral

ossification. Plays a role in wound healing, probably by a mechanism that involves proteolytic activation of TGFB1 and degradation of CCN2. Plays a role in keratinocyte migration during wound healing. May play a role in cell migration and in tumor cell invasion.

**Cellular Location**

Secreted, extracellular space, extracellular matrix. Secreted

**Tissue Location**

Detected in fetal cartilage and calvaria, in chondrocytes of hypertrophic cartilage in vertebrae and in the dorsal end of ribs undergoing ossification, as well as in osteoblasts and periosteal cells below the inner periosteal region of ossified ribs Detected in chondrocytes from in joint cartilage that have been treated with TNF and IL1B, but not in untreated chondrocytes. Detected in T lymphocytes. Detected in breast carcinoma tissue

**MMP13 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**MMP13 Antibody - Images****MMP13 Antibody - Background**

Degrades collagen type I. Does not act on gelatin or casein. Could have a role in tumoral process.

**MMP13 Antibody - References**

Freije J.M.P.,et al.J. Biol. Chem. 269:16766-16773(1994).  
Willmroth F.,et al.Immunobiology 198:375-384(1998).  
Gomis-Rueth F.-X.,et al.J. Mol. Biol. 264:556-566(1996).  
Kennedy A.M.,et al.J. Clin. Invest. 115:2832-2842(2005).  
Lausch E.,et al.Am. J. Hum. Genet. 85:168-178(2009).

**MMP13 Antibody - Citations**

- [Nkx2-5 Is Expressed in Atherosclerotic Plaques and Attenuates Development of Atherosclerosis in Apolipoprotein E-Deficient Mice.](#)