

PSMD12 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51456

Specification

PSMD12 Antibody - Product Information

Application WB
Primary Accession O00232

Reactivity Human, Mouse, Rat Rabbit

Clonality Polyclonal
Calculated MW 53 KDa
Antigen Region 131 - 190

PSMD12 Antibody - Additional Information

Gene ID 5718

Other Names

26S proteasome non-ATPase regulatory subunit 12, 26S proteasome regulatory subunit RPN5, 26S proteasome regulatory subunit p55, PSMD12

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human PSMD12. The exact sequence is proprietary.

Dilution

WB~~ 1:1000

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

PSMD12 Antibody - Protein Information

Name PSMD12

Function

Component of the 26S proteasome, a multiprotein complex involved in the ATP-dependent degradation of ubiquitinated proteins. This complex plays a key role in the maintenance of protein homeostasis by removing misfolded or damaged proteins, which could impair cellular functions, and by removing proteins whose functions are no longer required. Therefore, the proteasome participates in numerous cellular processes, including cell cycle progression, apoptosis, or DNA damage repair.




PSMD12 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

PSMD12 Antibody - Images



All lanes : Anti-PSMD12 Antibody at 1:1000 dilution Lane 1: HepG2 whole cell lysates Lane 2: U-87 MG whole cell lysates Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 53 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

PSMD12 Antibody - Background

Acts as a regulatory subunit of the 26S proteasome which is involved in the ATP-dependent degradation of ubiquitinated proteins.

PSMD12 Antibody - References

Saito A., et al. Gene 203:241-250(1997).

Ota T., et al. Nat. Genet. 36:40-45(2004).

Suzuki Y., et al. Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.

Zody M.C., et al. Nature 440:1045-1049(2006).

Mural R.J., et al. Submitted (JUL-2005) to the EMBL/GenBank/DDBJ databases.