

## SMG7 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51529

### Specification

# SMG7 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IP, ICC, IHC-P, E <u>092540</u> Human Rabbit Polyclonal 127 KDa

## SMG7 Antibody - Additional Information

Gene ID 9887

Other Names Protein SMG7, EST1-like protein C, SMG-7 homolog, hSMG-7, SMG7, C1orf16, EST1C, KIAA0250

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

# SMG7 Antibody - Protein Information

Name SMG7 (HGNC:16792)

Function

Plays a role in nonsense-mediated mRNA decay. Recruits UPF1 to cytoplasmic mRNA decay bodies. Together with SMG5 is thought to provide a link to the mRNA degradation machinery involving exonucleolytic pathways, and to serve as an adapter for UPF1 to protein phosphatase 2A (PP2A), thereby triggering UPF1 dephosphorylation.

Cellular Location

Cytoplasm. Nucleus. Note=Predominantly cytoplasmic, and nuclear. Shuttles between nucleus and cytoplasm

#### SMG7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides



- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety

<u>Cell Culture</u>

SMG7 Antibody - Images

## SMG7 Antibody - Background

Plays a role in nonsense-mediated mRNA decay. Recruits UPF1 to cytoplasmic mRNA decay bodies. Together with SMG5 is thought to provide a link to the mRNA degradation machinery involving exonucleolytic pathways, and to serve as an adapter for UPF1 to protein phosphatase 2A (PP2A), thereby triggering UPF1 dephosphorylation.

### SMG7 Antibody - References

Ohnishi T., et al.Mol. Cell 12:1187-1200(2003). Nagase T., et al.DNA Res. 3:321-329(1996). Nakajima D., et al.DNA Res. 9:99-106(2002). Ota T., et al.Nat. Genet. 36:40-45(2004). Gregory S.G., et al.Nature 441:315-321(2006).