

SRY Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51535

Specification

SRY Antibody - Product Information

Application	WB
Primary Accession	Q05066
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	24 KDa
Antigen Region	31 - 90

SRY Antibody - Additional Information

Gene ID 6736

Other Names

Sex-determining region Y protein, Testis-determining factor, SRY, TDF

Target/Specificity

KLH conjugated synthetic peptide derived from human SRY

Dilution

WB ~ 1:1000

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

SRY Antibody - Protein Information

Name SRY {ECO:0000303|PubMed:1695712, ECO:0000312|HGNC:HGNC:11311}

Function

Transcriptional regulator that controls a genetic switch in male development (PubMed:11563911). It is necessary and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells (PubMed:16414182, PubMed:16996051). Involved in different aspects of gene regulation including promoter activation or repression (PubMed:9525897). Binds to the DNA consensus sequence 5'- [AT]AACAA[AT]-3' (PubMed:11563911, PubMed:>1425584, PubMed:>15170344, PubMed:>8159753, PubMed:>8265659). SRY HMG box recognizes DNA by partial intercalation in the minor groove and promotes DNA bending (PubMed:>11563911, PubMed:>1425584, PubMed:>15170344, PubMed:>16762365, PubMed:>8159753, PubMed:>8265659). Also involved in pre-mRNA splicing (PubMed:>11818535). In male adult brain involved in the maintenance of motor functions of dopaminergic neurons (By similarity).

Cellular Location


Nucleus speckle. Cytoplasm Nucleus. Note=Acetylation contributes to its nuclear localization and deacetylation by HDAC3 induces a cytoplasmic delocalization (PubMed:15297880). Colocalizes with SOX6 in speckles (PubMed:11818535). Colocalizes with CAML in the nucleus (PubMed:15746192). Colocalizes in the nucleus with ZNF208 isoform KRAB- O and tyrosine hydroxylase (TH) (By similarity). Nuclear import is facilitated by XPO4, a protein that usually acts as a nuclear export signal receptor (PubMed:19349578). {ECO:0000250|UniProtKB:Q05738, ECO:0000269|PubMed:11818535, ECO:0000269|PubMed:15297880, ECO:0000269|PubMed:15746192, ECO:0000269|PubMed:19349578}

SRY Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SRY Antibody - Images



Anti-SRY Antibody at 1:1000 dilution + human testis lysates Lysates/proteins at 20 µg per lane.
Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

SRY Antibody - Background

Transcriptional regulator that controls a genetic switch in male development. It is necessary and sufficient for initiating male sex determination by directing the development of supporting cell precursors (pre-Sertoli cells) as Sertoli rather than granulosa cells (By similarity). In male adult brain involved in the maintenance of motor functions of dopaminergic neurons (By similarity). Involved in different aspects of gene regulation including promoter activation or repression (By similarity). Promotes DNA bending. SRY HMG box recognizes DNA by partial intercalation in the minor groove. Also involved in pre-mRNA splicing. Binds to the DNA consensus sequence 5'-[AT]AACAA[AT]-3'.

SRY Antibody - References

- Sinclair A.H.,et al.Nature 346:240-244(1990).
Su H.,et al.Am. J. Hum. Genet. 52:24-38(1993).
Behlke M.A.,et al.Genomics 17:736-739(1993).
Whitfield L.S.,et al.Genomics 27:306-311(1995).
Ferrari S.,et al.EMBO J. 11:4497-4506(1992).