

TRIP13 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51588

Specification

TRIP13 Antibody - Product Information

Application	WB, IHC-P, E
Primary Accession	Q15645
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	49 KDa

TRIP13 Antibody - Additional Information

Gene ID 9319

Other Names

Pachytene checkpoint protein 2 homolog, Human papillomavirus type 16 E1 protein-binding protein, 16E1-BP, HPV16 E1 protein-binding protein, Thyroid hormone receptor interactor 13, Thyroid receptor-interacting protein 13, TR-interacting protein 13, TRIP-13, TRIP13, PCH2

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the C-term region of human TRIP13. The exact sequence is proprietary.

Dilution

WB~~1:1000
IHC-P~~N/A
E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

TRIP13 Antibody - Protein Information

Name TRIP13

Synonyms PCH2

Function

Plays a key role in chromosome recombination and chromosome structure development during meiosis. Required at early steps in meiotic recombination that leads to non-crossovers pathways. Also needed for efficient completion of homologous synapsis by influencing crossover distribution along the chromosomes affecting both crossovers and non-crossovers pathways. Also required for

development of higher- order chromosome structures and is needed for synaptonemal-complex formation. In males, required for efficient synapsis of the sex chromosomes and for sex body formation. Promotes early steps of the DNA double-strand breaks (DSBs) repair process upstream of the assembly of RAD51 complexes. Required for depletion of HORMAD1 and HORMAD2 from synapsed chromosomes (By similarity). Plays a role in mitotic spindle assembly checkpoint (SAC) activation (PubMed:28553959).

TRIP13 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TRIP13 Antibody - Images

TRIP13 Antibody - Background

Plays a key role in chromosome recombination and chromosome structure development during meiosis. Required at early steps in meiotic recombination that leads to non-crossovers pathways. Also needed for efficient completion of homologous synapsis by influencing crossover distribution along the chromosomes affecting both crossovers and non-crossovers pathways. Also required for development of higher-order chromosome structures and is needed for synaptonemal-complex formation. In males, required for efficient synapsis of the sex chromosomes and for sex body formation. Promotes early steps of the DNA double- strand breaks (DSBs) repair process upstream of the assembly of RAD51 complexes. Required for depletion of HORMAD1 and HORMAD2 from synapsed chromosomes (By similarity).

TRIP13 Antibody - References

Yasugi T.,et al.J. Virol. 71:5942-5951(1997).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.
Lee J.W.,et al.Mol. Endocrinol. 9:243-254(1995).
Rush J.,et al.Nat. Biotechnol. 23:94-101(2005).