

**Ku70 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51617**

**Specification**

**Ku70 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P12956</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	70 KDa
Antigen Region	1 - 60

**Ku70 Antibody - Additional Information**

**Gene ID** 2547

**Other Names**

X-ray repair cross-complementing protein 6, 364-, 4299-, 5'-deoxyribose-5-phosphate lyase Ku70, 5'-dRP lyase Ku70, 70 kDa subunit of Ku antigen, ATP-dependent DNA helicase 2 subunit 1, ATP-dependent DNA helicase II 70 kDa subunit, CTC box-binding factor 75 kDa subunit, CTC75, CTCBF, DNA repair protein XRCC6, Lupus Ku autoantigen protein p70, Ku70, Thyroid-lupus autoantigen, TLAA, X-ray repair complementing defective repair in Chinese hamster cells 6, XRCC6, G22P1

**Target/Specificity**

KLH conjugated synthetic peptide derived from human Ku70

**Dilution**

WB~~ 1:1000

**Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**Ku70 Antibody - Protein Information**

**Name** XRCC6

**Synonyms** G22P1

**Function**

Single-stranded DNA-dependent ATP-dependent helicase that plays a key role in DNA non-homologous end joining (NHEJ) by recruiting DNA-PK to DNA (PubMed:[11493912](http://www.uniprot.org/citations/11493912)), PubMed:[11493912](http://www.ncbi.nlm.nih.gov/pubmed/11493912))

Required for double-strand break repair and V(D)J recombination (PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Required for double-strand break repair and V(D)J recombination (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Also has a role in chromosome translocation (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Has a role in chromosome translocation (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). It works in the 3'-5' direction (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). During NHEJ, the XRCC5-XRRC6 dimer performs the recognition step: it recognizes and binds to the broken ends of the DNA and protects them from further resection (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">9742108</a>). Binding to DNA may be mediated by XRCC6 (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">20493174</a>).

target="\_blank">>20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">>2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">>7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">>8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">>9742108</a>). The XRCC5-XRRC6 dimer acts as a regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">>11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">>12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">>20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">>2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">>7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">>8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">>9742108</a>). The XRCC5-XRRC6 dimer is probably involved in stabilizing broken DNA ends and bringing them together (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">>11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">>12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">>20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">>2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">>7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">>8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">>9742108</a>). The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step (PubMed:<a href="http://www.uniprot.org/citations/11493912" target="\_blank">>11493912</a>, PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">>12145306</a>, PubMed:<a href="http://www.uniprot.org/citations/20493174" target="\_blank">>20493174</a>, PubMed:<a href="http://www.uniprot.org/citations/2466842" target="\_blank">>2466842</a>, PubMed:<a href="http://www.uniprot.org/citations/7957065" target="\_blank">>7957065</a>, PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">>8621488</a>, PubMed:<a href="http://www.uniprot.org/citations/9742108" target="\_blank">>9742108</a>). Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose-5-phosphate at an abasic site near double-strand breaks (PubMed:<a href="http://www.uniprot.org/citations/20383123" target="\_blank">>20383123</a>). 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined (PubMed:<a href="http://www.uniprot.org/citations/20383123" target="\_blank">>20383123</a>). The XRCC5-XRRC6 dimer together with APEX1 acts as a negative regulator of transcription (PubMed:<a href="http://www.uniprot.org/citations/8621488" target="\_blank">>8621488</a>). In association with NAA15, the XRCC5-XRRC6 dimer binds to the osteocalcin promoter and activates osteocalcin expression (PubMed:<a href="http://www.uniprot.org/citations/12145306" target="\_blank">>12145306</a>). Plays a role in the regulation of DNA virus-mediated innate immune response by assembling into the HDP-RNP complex, a complex that serves as a platform for IRF3 phosphorylation and subsequent innate immune response activation through the cGAS-STING pathway (PubMed:<a href="http://www.uniprot.org/citations/28712728" target="\_blank">>28712728</a>). Negatively regulates apoptosis by interacting with BAX and sequestering it from the mitochondria (PubMed:<a href="http://www.uniprot.org/citations/15023334" target="\_blank">>15023334</a>). Might have deubiquitination activity, acting on BAX (PubMed:<a href="http://www.uniprot.org/citations/18362350" target="\_blank">>18362350</a>).

### Cellular Location


Nucleus. Chromosome. Cytoplasm. Note=When trimethylated, localizes in the cytoplasm.

### Ku70 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### Ku70 Antibody - Images



All lanes : Anti-Ku70 Antibody at 1:1000 dilution Lane 1: HeLa whole cell lysates Lane 2: HCT116 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 70 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### Ku70 Antibody - Background

Single-stranded DNA-dependent ATP-dependent helicase. Has a role in chromosome translocation. The DNA helicase II complex binds preferentially to fork-like ends of double-stranded DNA in a cell cycle-dependent manner. It works in the 3'-5' direction. Binding to DNA may be mediated by XRCC6. Involved in DNA non-homologous end joining (NHEJ) required for double-strand break repair and V(D)J recombination. The XRCC5/6 dimer acts as regulatory subunit of the DNA-dependent protein kinase complex DNA-PK by increasing the affinity of the catalytic subunit PRKDC to DNA by 100-fold. The XRCC5/6 dimer is probably involved in stabilizing broken DNA ends and bringing them together. The assembly of the DNA-PK complex to DNA ends is required for the NHEJ ligation step. Required for osteocalcin gene expression. Probably also acts as a 5'-deoxyribose-5-phosphate lyase (5'-dRP lyase), by catalyzing the beta-elimination of the 5' deoxyribose- 5-phosphate at an abasic site near double-strand breaks. 5'-dRP lyase activity allows to 'clean' the termini of abasic sites, a class of nucleotide damage commonly associated with strand breaks, before such broken ends can be joined. The XRCC5/6 dimer together with APEX1 acts as a negative regulator of transcription.

### Ku70 Antibody - References

- Chan J.Y., et al. J. Biol. Chem. 264:3651-3654(1989).  
Reeves W.H., et al. J. Biol. Chem. 264:5047-5052(1989).  
Griffith A.J., et al. Mol. Biol. Rep. 16:91-97(1992).  
Halleck A., et al. Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

Dunham I.,et al.Nature 402:489-495(1999).