

TTPAL Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5164b**Specification**

TTPAL Antibody (C-term) - Product Information

Application	FC, IHC-P, WB,E
Primary Accession	Q9BTX7
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	38515
Antigen Region	204-233

TTPAL Antibody (C-term) - Additional Information**Gene ID** 79183**Other Names**

Alpha-tocopherol transfer protein-like, TTPAL, C20orf121

Target/Specificity

This TTPAL antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 204-233 amino acids from the C-terminal region of human TTPAL.

Dilution

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

TTPAL Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

TTPAL Antibody (C-term) - Protein Information**Name** TTPAL

Synonyms C20orf121


Function May act as a protein that binds a hydrophobic ligand.

TTPAL Antibody (C-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


TTPAL Antibody (C-term) - Images



Western blot analysis of TTPAL Antibody (C-term) (Cat. #AP5164b) in MCF-7 cell line lysates (35ug/lane). TTPAL (arrow) was detected using the purified Pab.



TTPAL Antibody (C-term) (Cat. #AP5164b) IHC analysis in formalin fixed and paraffin embedded human breast carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the TTPAL Antibody (C-term) for immunohistochemistry. Clinical relevance has not been evaluated.



TTPAL Antibody (C-term) (Cat. #AP5164b) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

TTPAL Antibody (C-term) - Background

TTPAL may act as a protein that binds a hydrophobic ligand.

TTPAL Antibody (C-term) - References

Deloukas, P., et al. Nature 414(6866):865-871(2001)