

**RPS3 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51707**

**Specification**

---

**RPS3 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P23396</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	33 KDa
Antigen Region	161 - 220

**RPS3 Antibody - Additional Information**

**Gene ID** 6188

**Other Names**

40S ribosomal protein S3, RPS3

**Target/Specificity**

KLH conjugated synthetic peptide derived from human RPS3

**Dilution**

WB~~ 1:1000

**Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**RPS3 Antibody - Protein Information**

**Name** RPS3 {ECO:0000303|PubMed:11875025}

**Function**

Component of the small ribosomal subunit (PubMed:<a href="http://www.uniprot.org/citations/8706699" target="\_blank">8706699</a>, PubMed:<a href="http://www.uniprot.org/citations/23636399" target="\_blank">23636399</a>). The ribosome is a large ribonucleoprotein complex responsible for the synthesis of proteins in the cell (PubMed:<a href="http://www.uniprot.org/citations/8706699" target="\_blank">8706699</a>, PubMed:<a href="http://www.uniprot.org/citations/23636399" target="\_blank">23636399</a>). Has endonuclease activity and plays a role in repair of damaged DNA (PubMed:<a href="http://www.uniprot.org/citations/7775413" target="\_blank">7775413</a>). Cleaves phosphodiester bonds of DNAs containing altered bases with broad specificity and cleaves supercoiled DNA more efficiently than relaxed DNA (PubMed:<a

href="http://www.uniprot.org/citations/15707971" target="\_blank">15707971</a>). Displays high binding affinity for 7,8-dihydro-8-oxoguanine (8-oxoG), a common DNA lesion caused by reactive oxygen species (ROS) (PubMed:<a href="http://www.uniprot.org/citations/14706345" target="\_blank">14706345</a>). Has also been shown to bind with similar affinity to intact and damaged DNA (PubMed:<a href="http://www.uniprot.org/citations/18610840" target="\_blank">18610840</a>). Stimulates the N-glycosylase activity of the base excision protein OGG1 (PubMed:<a href="http://www.uniprot.org/citations/15518571" target="\_blank">15518571</a>). Enhances the uracil excision activity of UNG1 (PubMed:<a href="http://www.uniprot.org/citations/18973764" target="\_blank">18973764</a>). Also stimulates the cleavage of the phosphodiester backbone by APEX1 (PubMed:<a href="http://www.uniprot.org/citations/18973764" target="\_blank">18973764</a>). When located in the mitochondrion, reduces cellular ROS levels and mitochondrial DNA damage (PubMed:<a href="http://www.uniprot.org/citations/23911537" target="\_blank">23911537</a>). Has also been shown to negatively regulate DNA repair in cells exposed to hydrogen peroxide (PubMed:<a href="http://www.uniprot.org/citations/17049931" target="\_blank">17049931</a>). Plays a role in regulating transcription as part of the NF-kappa-B p65-p50 complex where it binds to the RELA/p65 subunit, enhances binding of the complex to DNA and promotes transcription of target genes (PubMed:<a href="http://www.uniprot.org/citations/18045535" target="\_blank">18045535</a>). Represses its own translation by binding to its cognate mRNA (PubMed:<a href="http://www.uniprot.org/citations/20217897" target="\_blank">20217897</a>). Binds to and protects TP53/p53 from MDM2-mediated ubiquitination (PubMed:<a href="http://www.uniprot.org/citations/19656744" target="\_blank">19656744</a>). Involved in spindle formation and chromosome movement during mitosis by regulating microtubule polymerization (PubMed:<a href="http://www.uniprot.org/citations/23131551" target="\_blank">23131551</a>). Involved in induction of apoptosis through its role in activation of CASP8 (PubMed:<a href="http://www.uniprot.org/citations/14988002" target="\_blank">14988002</a>). Induces neuronal apoptosis by interacting with the E2F1 transcription factor and acting synergistically with it to up-regulate pro-apoptotic proteins BCL2L11/BIM and HRK/Dp5 (PubMed:<a href="http://www.uniprot.org/citations/20605787" target="\_blank">20605787</a>). Interacts with TRADD following exposure to UV radiation and induces apoptosis by caspase-dependent JNK activation (PubMed:<a href="http://www.uniprot.org/citations/22510408" target="\_blank">22510408</a>).

### **Cellular Location**

Cytoplasm. Nucleus. Nucleus, nucleolus Mitochondrion inner membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton, spindle. Note=In normal cells, located mainly in the cytoplasm with small amounts in the nucleus but translocates to the nucleus in cells undergoing apoptosis (By similarity). Nuclear translocation is induced by DNA damaging agents such as hydrogen peroxide (PubMed:17560175). Accumulates in the mitochondrion in response to increased ROS levels (PubMed:23911537) Localizes to the spindle during mitosis (PubMed:23131551). Localized in cytoplasmic mRNP granules containing untranslated mRNAs (PubMed:17289661).


{ECO:0000250|UniProtKB:P62908, ECO:0000269|PubMed:17289661,  
ECO:0000269|PubMed:17560175, ECO:0000269|PubMed:23131551,  
ECO:0000269|PubMed:23911537}

### **RPS3 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)

- [Flow Cytometry](#)
- [Cell Culture](#)

**RPS3 Antibody - Images**

All lanes : Anti-RPS3 Antibody at 1:1000 dilution Lane 1: Hela whole cell lysates Lane 2: NIH/3T3 whole cell lysates Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution Predicted band size : 27 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

**RPS3 Antibody - References**

- Zhang X.T.,et al.Nucleic Acids Res. 18:6689-6689(1990).  
Pogue-Geile K.,et al.Mol. Cell. Biol. 11:3842-3849(1991).  
Yoshihama M.,et al.Genome Res. 12:379-390(2002).  
Ota T.,et al.Nat. Genet. 36:40-45(2004).  
Taylor T.D.,et al.Nature 440:497-500(2006).