

**AP2M1 Antibody**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP51726**

**Specification**

---

**AP2M1 Antibody - Product Information**

Application	WB, E
Primary Accession	<a href="#">Q96CW1</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	50 KDa

**AP2M1 Antibody - Additional Information**

**Gene ID 1173**

**Other Names**

AP-2 complex subunit mu, AP-2 mu chain, Adapton-mu2, Adaptor protein complex AP-2 subunit mu, Adaptor-related protein complex 2 subunit mu, Clathrin assembly protein complex 2 mu medium chain, Clathrin coat assembly protein AP50, Clathrin coat-associated protein AP50, HA2 50 kDa subunit, Plasma membrane adaptor AP-2 50 kDa protein, AP2M1, CLAPM1, KIAA0109

**Dilution**

WB~~1:1000  
E~~N/A

**Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

**Storage**

Store at -20 °C. Stable for 12 months from date of receipt

**AP2M1 Antibody - Protein Information**

**Name** [AP2M1 \(HGNC:564\)](#)

**Synonyms** CLAPM1, KIAA0109

**Function**

Component of the adaptor protein complex 2 (AP-2) (PubMed:[12694563](http://www.uniprot.org/citations/12694563), PubMed:[12952941](http://www.uniprot.org/citations/12952941), PubMed:[14745134](http://www.uniprot.org/citations/14745134), PubMed:[14985334](http://www.uniprot.org/citations/14985334), PubMed:[15473838](http://www.uniprot.org/citations/15473838), PubMed:[31104773](http://www.uniprot.org/citations/31104773)). Adaptor protein complexes function in protein transport via transport vesicles in different membrane traffic

pathways (PubMed:<a href="http://www.uniprot.org/citations/12694563" target="\_blank">12694563</a>, PubMed:<a href="http://www.uniprot.org/citations/12952941" target="\_blank">12952941</a>, PubMed:<a href="http://www.uniprot.org/citations/14745134" target="\_blank">14745134</a>, PubMed:<a href="http://www.uniprot.org/citations/14985334" target="\_blank">14985334</a>, PubMed:<a href="http://www.uniprot.org/citations/15473838" target="\_blank">15473838</a>, PubMed:<a href="http://www.uniprot.org/citations/31104773" target="\_blank">31104773</a>). Adaptor protein complexes are vesicle coat components and appear to be involved in cargo selection and vesicle formation (PubMed:<a href="http://www.uniprot.org/citations/12694563" target="\_blank">12694563</a>, PubMed:<a href="http://www.uniprot.org/citations/12952941" target="\_blank">12952941</a>, PubMed:<a href="http://www.uniprot.org/citations/14745134" target="\_blank">14745134</a>, PubMed:<a href="http://www.uniprot.org/citations/14985334" target="\_blank">14985334</a>, PubMed:<a href="http://www.uniprot.org/citations/15473838" target="\_blank">15473838</a>, PubMed:<a href="http://www.uniprot.org/citations/31104773" target="\_blank">31104773</a>). AP-2 is involved in clathrin-dependent endocytosis in which cargo proteins are incorporated into vesicles surrounded by clathrin (clathrin-coated vesicles, CCVs) which are destined for fusion with the early endosome (PubMed:<a href="http://www.uniprot.org/citations/12694563" target="\_blank">12694563</a>, PubMed:<a href="http://www.uniprot.org/citations/12952941" target="\_blank">12952941</a>, PubMed:<a href="http://www.uniprot.org/citations/14745134" target="\_blank">14745134</a>, PubMed:<a href="http://www.uniprot.org/citations/14985334" target="\_blank">14985334</a>, PubMed:<a href="http://www.uniprot.org/citations/15473838" target="\_blank">15473838</a>, PubMed:<a href="http://www.uniprot.org/citations/31104773" target="\_blank">31104773</a>). AP-2 is unable to bind directly to membrane components (PubMed:<a href="http://www.uniprot.org/citations/12694563" target="\_blank">12694563</a>, PubMed:<a href="http://www.uniprot.org/citations/12952941" target="\_blank">12952941</a>, PubMed:<a href="http://www.uniprot.org/citations/14745134" target="\_blank">14745134</a>, PubMed:<a href="http://www.uniprot.org/citations/14985334" target="\_blank">14985334</a>, PubMed:<a href="http://www.uniprot.org/citations/15473838" target="\_blank">15473838</a>, PubMed:<a href="http://www.uniprot.org/citations/31104773" target="\_blank">31104773</a>).

Clathrin-associated adaptor protein (AP) complexes which can bind directly to both the clathrin lattice and to the lipid and protein components of membranes are considered to be the major clathrin adaptors contributing the CCV formation (PubMed:<a href="http://www.uniprot.org/citations/12694563" target="\_blank">12694563</a>, PubMed:<a href="http://www.uniprot.org/citations/12952941" target="\_blank">12952941</a>, PubMed:<a href="http://www.uniprot.org/citations/14745134" target="\_blank">14745134</a>, PubMed:<a href="http://www.uniprot.org/citations/14985334" target="\_blank">14985334</a>, PubMed:<a href="http://www.uniprot.org/citations/15473838" target="\_blank">15473838</a>, PubMed:<a href="http://www.uniprot.org/citations/31104773" target="\_blank">31104773</a>). AP-2 also serves as a cargo receptor to selectively sort the membrane proteins involved in receptor-mediated endocytosis (PubMed:<a href="http://www.uniprot.org/citations/16581796" target="\_blank">16581796</a>). AP-2 seems to play a role in the recycling of synaptic vesicle membranes from the presynaptic surface (PubMed:<a href="http://www.uniprot.org/citations/12694563" target="\_blank">12694563</a>, PubMed:<a href="http://www.uniprot.org/citations/12952941" target="\_blank">12952941</a>, PubMed:<a href="http://www.uniprot.org/citations/14745134" target="\_blank">14745134</a>, PubMed:<a href="http://www.uniprot.org/citations/14985334" target="\_blank">14985334</a>, PubMed:<a href="http://www.uniprot.org/citations/15473838" target="\_blank">15473838</a>, PubMed:<a href="http://www.uniprot.org/citations/31104773" target="\_blank">31104773</a>). AP-2 recognizes Y-X-X-[FILMV] (Y-X- X-Phi) and [ED]-X-X-X-L-[LI] endocytosis signal motifs within the cytosolic tails of transmembrane cargo molecules (By similarity). AP-2 may also play a role in maintaining normal post-endocytic trafficking through the ARF6-regulated, non-clathrin pathway (PubMed:<a href="http://www.uniprot.org/citations/19033387" target="\_blank">19033387</a>). During long-term potentiation in hippocampal neurons, AP-2 is responsible for the endocytosis of ADAM10 (PubMed:<a href="http://www.uniprot.org/citations/23676497" target="\_blank">23676497</a>). The AP-2 mu subunit binds to transmembrane cargo proteins; it recognizes the Y- X-X-Phi motifs (By similarity). The surface region interacting with to the Y-X-X-Phi

motif is inaccessible in cytosolic AP-2, but becomes accessible through a conformational change following phosphorylation of AP-2 mu subunit at Thr-156 in membrane-associated AP-2 (PubMed:<a href="http://www.uniprot.org/citations/11877457" target="\_blank">11877457</a>). The membrane-specific phosphorylation event appears to involve assembled clathrin which activates the AP-2 mu kinase AAK1 (PubMed:<a href="http://www.uniprot.org/citations/11877457" target="\_blank">11877457</a>). Plays a role in endocytosis of frizzled family members upon Wnt signaling (By similarity).

#### **Cellular Location**

Cell membrane. Membrane, coated pit; Peripheral membrane protein; Cytoplasmic side.  
Note=AP-2 appears to be excluded from internalizing CCVs and to disengage from sites of endocytosis seconds before internalization of the nascent CCV {ECO:0000250|UniProtKB:P84091}

#### **Tissue Location**

Expressed in the brain (at protein level).

### **AP2M1 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **AP2M1 Antibody - Images**

### **AP2M1 Antibody - Background**

Component of the adaptor protein complex 2 (AP-2). Adaptor protein complexes function in protein transport via transport vesicles in different membrane traffic pathways. Adaptor protein complexes are vesicle coat components and appear to be involved in cargo selection and vesicle formation. AP-2 is involved in clathrin-dependent endocytosis in which cargo proteins are incorporated into vesicles surrounded by clathrin (clathrin- coated vesicles, CCVs) which are destined for fusion with the early endosome. The clathrin lattice serves as a mechanical scaffold but is itself unable to bind directly to membrane components. Clathrin-associated adaptor protein (AP) complexes which can bind directly to both the clathrin lattice and to the lipid and protein components of membranes are considered to be the major clathrin adaptors contributing the CCV formation. AP-2 also serves as a cargo receptor to selectively sort the membrane proteins involved in receptor-mediated endocytosis. AP-2 seems to play a role in the recycling of synaptic vesicle membranes from the presynaptic surface. AP-2 recognizes Y-X-X-[FILMV] (Y-X-X-Phi) and [ED]-X-X-X-L-[LI] endocytosis signal motifs within the cytosolic tails of transmembrane cargo molecules. AP-2 may also play a role in maintaining normal post-endocytic trafficking through the ARF6-regulated, non-clathrin pathway. The AP-2 mu subunit binds to transmembrane cargo proteins; it recognizes the Y-X-X-Phi motifs. The surface region interacting with to the Y-X-X-Phi motif is inaccessible in cytosolic AP-2, but becomes accessible through a conformational change following phosphorylation of AP-2 mu subunit at 'Tyr-156' in membrane- associated AP-2. The membrane-specific phosphorylation event appears to involve assembled clathrin which activates the AP-2 mu kinase AAK1 (By similarity). Plays a role in endocytosis of frizzled family members upon Wnt signaling (By similarity).

### **AP2M1 Antibody - References**

Tsui S.K.W.,et al.Submitted (APR-1996) to the EMBL/GenBank/DDBJ databases.  
Nagase T.,et al.DNA Res. 2:37-43(1995).  
Kalinine N.,et al.Submitted (MAY-2003) to the EMBL/GenBank/DDBJ databases.  
Muzny D.M.,et al.Nature 440:1194-1198(2006).  
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.