

DHRS2 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51756**Specification**

DHRS2 Antibody - Product Information

Application	WB, E
Primary Accession	Q13268
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	27 KDa

DHRS2 Antibody - Additional Information**Gene ID** 10202**Other Names**

Dehydrogenase/reductase SDR family member 2, mitochondrial, 111-, Dicarbonyl reductase HEP27, Protein D, DHRS2

Dilution

WB~~1:1000

E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C.Stable for 12 months from date of receipt

DHRS2 Antibody - Protein Information**Name** DHRS2 ([HGNC:18349](#))**Synonyms** SDR25C1**Function**

NADPH-dependent oxidoreductase which catalyzes the reduction of dicarbonyl compounds. Displays reductase activity in vitro with 3,4- hexanedione, 2,3-heptanedione and 1-phenyl-1,2-propanedione as substrates (PubMed:16685466). May function as a dicarbonyl reductase in the enzymatic inactivation of reactive carbonyls involved in covalent modification of cellular components (PubMed:16685466). Also displays a minor hydroxysteroid dehydrogenase activity toward bile acids such as ursodeoxycholic acid (UDCA) and isoursodeoxycholic acid (isoUDCA), which makes it unlikely to control hormone levels (PubMed:16685466).

Doesn't show any activity in vitro with retinoids and sugars as substrates (PubMed:16685466). Attenuates MDM2-mediated p53/TP53 degradation, leading to p53/TP53 stabilization and increased transcription activity, resulting in the accumulation of MDM2 and CDKN1A/p21 (PubMed:20547751). Reduces proliferation, migration and invasion of cancer cells and well as the production of ROS in cancer (PubMed:29106393).

Cellular Location

Mitochondrion matrix. Nucleus. Note=A minor fraction of the protein is translocated from the mitochondria to the nucleus, after cleavage of the targeting signal

Tissue Location

Widely expressed, with highest levels in liver and kidney, followed by heart, spleen, skeletal muscle and placenta. In hemopoietic cells, expressed in dendritic cells, but not in monocytes, macrophages, granulocytes, nor in B and T lymphocytes

DHRS2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DHRS2 Antibody - Images

DHRS2 Antibody - Background

Displays NADPH-dependent dicarbonyl reductase activity in vitro with 3,4-Hexanedione, 2,3-Heptanedione and 1-Phenyl-1,2- propanedione as substrates. No reductase activity is displayed in vitro with steroids, retinoids and sugars as substrates. Attenuates MDM2-mediated p53/TP53 degradation, leading to p53/TP53 stabilization and increased transcription activity, resulting in the accumulation of MDM2 and CDKN1A/p21.

DHRS2 Antibody - References

Gabrielli F.,et al.Eur. J. Biochem. 232:473-477(1995).
Pellegrini S.,et al.Biochim. Biophys. Acta 1574:215-222(2002).
Suzuki Y.,et al.Submitted (APR-2005) to the EMBL/GenBank/DDBJ databases.
Heilig R.,et al.Nature 421:601-607(2003).
Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.