

CD248 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51768

Specification

CD248 Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, E
O9HCU0
Human, Mouse, Rat
Rabbit
Polyclonal
80 KDa

CD248 Antibody - Additional Information

Gene ID 57124

Other Names

Endosialin, Tumor endothelial marker 1, CD248, CD248, CD164L1, TEM1

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD248. The exact sequence is proprietary.

Dilution

WB~~1:1000 E~~N/A

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

CD248 Antibody - Protein Information

Name CD248

Synonyms CD164L1, TEM1

Function

Cell surface glycoprotein involved in various biological processes including angiogenesis, immune response modulation, and tissue remodeling and repair. Participates in pericyte proliferation through positive modulation of the PDGF receptor signaling pathway (PubMed:20484976). Acts as a scaffold for factor X, triggering allosteric changes and the spatial re-alignment of factor X with the TF-factor VIIa complex, thereby enhancing coagulation activation. Modulates the insulin signaling pathway by interacting with insulin receptor/INSR and by diminishing its capacity to be



autophosphorylated in response to insulin. Also regulates LPS-induced inflammatory response in macrophages by favoring the production of proinflammatory cytokines. In human, negatively regulates T-cell proliferation compared with stromal cells where it increases proliferation (PubMed:21466550).

Cellular Location

Membrane; Single-pass type I membrane protein

Tissue Location

Expressed in tumor endothelial cells but absent or barely detectable in normal endothelial cells. Expressed in metastatic lesions of the liver and during angiogenesis of corpus luteum formation and wound healing. Expressed in vascular endothelial cells of malignant tumors but not in normal blood vessels. Expressed in stromal fibroblasts. Strongly expressed in pericytes (PubMed:20484976) Expressed on stromal cells and cells with lymphoid morphology such a T- cells (PubMed:21466550).

CD248 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CD248 Antibody - Images

CD248 Antibody - Background

May play a role in tumor angiogenesis.

CD248 Antibody - References

St Croix B.,et al.Science 289:1197-1202(2000). Christian S.,et al.J. Biol. Chem. 276:7408-7414(2001). Ota T.,et al.Nat. Genet. 36:40-45(2004). Rettig W.J.,et al.Proc. Natl. Acad. Sci. U.S.A. 89:10832-10836(1992). Dolznig H.,et al.Cancer Immun. 5:10-10(2005).