

# **MASP2 Antibody**

Antigen-affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51788

# **Specification**

# **MASP2 Antibody - Product Information**

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW
Antigen Region

WB
000187
Human, Mouse
Rabbit
Polyclonal
76 KDa
211 - 270

# **MASP2 Antibody - Additional Information**

# **Gene ID 10747**

#### **Other Names**

Mannan-binding lectin serine protease 2, MBL-associated serine protease 2, Mannose-binding protein-associated serine protease 2, MASP-2, Mannan-binding lectin serine protease 2 A chain, Mannan-binding lectin serine protease 2 B chain, MASP2

### Target/Specificity

KLH conjugated synthetic peptide derived from human MASP2

#### **Dilution**

WB~~ 1:1000

#### **Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

### **MASP2 Antibody - Protein Information**

# Name MASP2

### **Function**

Serum protease that plays an important role in the activation of the complement system via mannose-binding lectin. After activation by auto-catalytic cleavage it cleaves C2 and C4, leading to their activation and to the formation of C3 convertase.

# **Cellular Location**

Secreted.

# **Tissue Location**




Plasma.

# **MASP2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# MASP2 Antibody - Images



Anti-MASP2 Antibody at 1:1000 dilution + HeLa whole cell lysates Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L),Peroxidase conjugated at 1/10000 dilution Predicted band size : 76 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### MASP2 Antibody - Background

Serum protease that plays an important role in the activation of the complement system via mannose-binding lectin. After activation by auto-catalytic cleavage it cleaves C2 and C4, leading to their activation and to the formation of C3 convertase.

# **MASP2 Antibody - References**

Thiel S., et al. Nature 386:506-510(1997).

Takahashi M., et al. Int. Immunol. 11:859-863(1999).

Stover C.M., et al.J. Immunol. 162:3481-3490(1999).

Thiel S., et al. Submitted (JUL-1998) to the EMBL/GenBank/DDBJ databases.

Takahashi M., et al. Submitted (OCT-1999) to the EMBL/GenBank/DDBJ databases.