

SLU7 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51824

Specification

SLU7 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, E <u>095391</u> Human, Mouse, Rat Rabbit Polyclonal 68 KDa

SLU7 Antibody - Additional Information

Gene ID 10569

Other Names Pre-mRNA-splicing factor SLU7, hSlu7, SLU7

Dilution WB~~1:1000 E~~N/A

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

SLU7 Antibody - Protein Information

Name SLU7

Function

Required for pre-mRNA splicing as component of the spliceosome (PubMed:10197984, PubMed:28502770, PubMed:30705154). Participates in the second catalytic step of pre-mRNA splicing, when the free hydroxyl group of exon I attacks the 3'-splice site to generate spliced mRNA and the excised lariat intron. Required for holding exon 1 properly in the spliceosome and for correct AG identification when more than one possible AG exists in 3'-splicing site region. May be involved in the activation of proximal AG. Probably also involved in alternative splicing regulation.

Cellular Location

Nucleus. Nucleus speckle. Cytoplasm Note=Predominantly nuclear. Shuttling between the nucleus and the cytoplasm is regulated by the CCHC-type zinc finger. Upon UV-C stress stimulus, the



nuclear concentration of the protein decreases, affecting alternative splicing. Translocates from the nucleus to the cytoplasm after heat shock cell treatment. Accumulates in cytoplasmic vesicle-like organelles after heat shock treatment, which may represent stress granules.

SLU7 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

SLU7 Antibody - Images

SLU7 Antibody - Background

Participates in the second catalytic step of pre-mRNA splicing, when the free hydroxyl group of exon I attacks the 3'- splice site to generate spliced mRNA and the excised lariat intron. Required for holding exon 1 properly in the spliceosome and for correct AG identification when more than one possible AG exists in 3'-splicing site region. May be involved in the activation of proximal AG. Probably also involved in alternative splicing regulation.

SLU7 Antibody - References

Chua K.,et al.Genes Dev. 13:841-850(1999). Schmutz J.,et al.Nature 431:268-274(2004). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Holste D.,et al.Submitted (AUG-2005) to the EMBL/GenBank/DDBJ databases. Chua K.,et al.Nature 402:207-210(1999).