

CDC42 Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51888

Specification

CDC42 Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, E
P60953
Human, Mouse, Rat
Rabbit
Polyclonal
21 KDa

CDC42 Antibody - Additional Information

Gene ID 998

Other Names

Cell division control protein 42 homolog, G25K GTP-binding protein, CDC42

Format

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage

Store at -20 °C. Stable for 12 months from date of receipt

CDC42 Antibody - Protein Information

Name CDC42 (HGNC:1736)

Function

Plasma membrane-associated small GTPase which cycles between an active GTP-bound and an inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses. Involved in epithelial cell polarization processes. Regulates the bipolar attachment of spindle microtubules to kinetochores before chromosome congression in metaphase (PubMed: 15642749). Regulates cell migration (PubMed: 17038317). In neurons, plays a role in the extension and maintenance of the formation of filopodia, thin and actin-rich surface projections (PubMed:14978216). Required for DOCK10-mediated spine formation in Purkinje cells and hippocampal neurons. In podocytes, facilitates filopodia and podosomes formation upon DOCK11- activation (PubMed: 33523862). Upon activation by CaMKII, modulates dendritic spine structural plasticity by relaying CaMKII transient activation to synapse-specific, long-term signaling (By similarity). Also plays a role in phagocytosis through organization of the F-actin cytoskeleton associated with forming phagocytic cups (PubMed:26465210).



Cellular Location

Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, spindle. Midbody Cell projection, dendrite {ECO:0000250|UniProtKB:P60766} Note=Localizes to spindle during prometaphase cells. Moves to the central spindle as cells progressed through anaphase to telophase (PubMed:15642749). Localizes at the end of cytokinesis in the intercellular bridge formed between two daughter cells (PubMed:15642749). Its localization is regulated by the activities of guanine nucleotide exchange factor ECT2 and GTPase activating protein RACGAP1 (PubMed:15642749). Colocalizes with NEK6 in the centrosome (PubMed:20873783). In its active GTP-bound form localizes to the leading edge membrane of migrating dendritic cells (By similarity) {ECO:0000250|UniProtKB:P60766, ECO:0000269|PubMed:15642749, ECO:0000269|PubMed:20873783}

CDC42 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CDC42 Antibody - Images

CDC42 Antibody - Background

Plasma membrane-associated small GTPase which cycles between an active GTP-bound and an inactive GDP-bound state. In active state binds to a variety of effector proteins to regulate cellular responses. Involved in epithelial cell polarization processes. Regulates the bipolar attachment of spindle microtubules to kinetochores before chromosome congression in metaphase. Plays a role in the extension and maintenance of the formation of thin, actin-rich surface projections called filopodia. Mediates CDC42-dependent cell migration.

CDC42 Antibody - References

Munemitsu S.,et al.Mol. Cell. Biol. 10:5977-5982(1990). Shinjo K.,et al.Proc. Natl. Acad. Sci. U.S.A. 87:9853-9857(1990). Rhodes S.,et al.Submitted (OCT-1999) to the EMBL/GenBank/DDBJ databases. Puhl H.L. III,et al.Submitted (APR-2002) to the EMBL/GenBank/DDBJ databases. Gregory S.G.,et al.Nature 441:315-321(2006).