

#### **PS2 Antibody**

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP51891

## **Specification**

## **PS2 Antibody - Product Information**

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, E
P04155
Human, Mouse, Rat
Rabbit
Polyclonal
11 KDa

## **PS2 Antibody - Additional Information**

**Gene ID 7031** 

#### **Other Names**

Trefoil factor 1, Breast cancer estrogen-inducible protein, PNR-2, Polypeptide P1A, hP1A, Protein pS2, TFF1, BCEI, PS2

## **Dilution**

WB~~1:1000 E~~N/A

#### **Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

## Storage

Store at -20 °C. Stable for 12 months from date of receipt

## **PS2 Antibody - Protein Information**

## Name TFF1

Synonyms BCEI, PS2

## **Function**

Stabilizer of the mucous gel overlying the gastrointestinal mucosa that provides a physical barrier against various noxious agents. May inhibit the growth of calcium oxalate crystals in urine.

#### **Cellular Location**

Secreted

## **Tissue Location**

Found in stomach, with highest levels in the upper gastric mucosal cells (at protein level). Detected in goblet cells of the small and large intestine and rectum, small submucosal glands in the esophagus, mucous acini of the sublingual gland, submucosal glands of the trachea, and



epithelial cells lining the exocrine pancreatic ducts but not in the remainder of the pancreas (at protein level) Scattered expression is detected in the epithelial cells of the gallbladder and submucosal glands of the vagina, and weak expression is observed in the bronchial goblet cells of the pseudostratified epithelia in the respiratory system (at protein level). Detected in urine (at protein level). Strongly expressed in breast cancer but at low levels in normal mammary tissue. It is regulated by estrogen in MCF-7 cells. Strong expression found in normal gastric mucosa and in the regenerative tissues surrounding ulcerous lesions of gastrointestinal tract, but lower expression found in gastric cancer (at protein level).

## **PS2 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **PS2 Antibody - Images**

## **PS2 Antibody - Background**

Stabilizer of the mucous gel overlying the gastrointestinal mucosa that provides a physical barrier against various noxious agents. May inhibit the growth of calcium oxalate crystals in urine.

# **PS2 Antibody - References**

Jakowlew S.B., et al. Nucleic Acids Res. 12:2861-2878(1984). Prud'Homme J.-F., et al. DNA 4:11-21(1985). Jeltsch J.-M., et al. Nucleic Acids Res. 15:1401-1414(1987). Takahashi H., et al. FEBS Lett. 261:283-286(1990). Mori K., et al. J. Biochem. 107:73-76(1990).