

## TIG2 Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP51981

## Specification

# TIG2 Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, E <u>O99969</u> Human, Mouse, Rat Rabbit Polyclonal 18 KDa

## TIG2 Antibody - Additional Information

Gene ID 5919

**Other Names** Retinoic acid receptor responder protein 2, Chemerin, RAR-responsive protein TIG2, Tazarotene-induced gene 2 protein, RARRES2, TIG2

**Dilution** WB~~1:1000 E~~N/A

Format 0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

Storage Store at -20 °C.Stable for 12 months from date of receipt

## **TIG2 Antibody - Protein Information**

Name RARRES2

Synonyms TIG2

### Function

Adipocyte-secreted protein (adipokine) that regulates adipogenesis, metabolism and inflammation through activation of the chemokine-like receptor 1 (CMKLR1). Also acts as a ligand for CMKLR2. Can also bind to C-C chemokine receptor-like 2 (CCRL2), but with a lower affinity than it does to CMKLR1 or CMKLR2 (PubMed:<a href="http://www.uniprot.org/citations/27716822" target="\_blank">27716822</a>). Positively regulates adipocyte differentiation, modulates the expression of adipocyte genes involved in lipid and glucose metabolism and might play a role in angiogenesis, a process essential for the expansion of white adipose tissue. Also acts as a pro-inflammatory adipokine, causing an increase in secretion of pro-inflammatory and prodiabetic adipokines, which further impair adipose tissue metabolic function and have negative systemic effects including impaired insulin sensitivity, altered glucose and lipid metabolism, and a decrease



in vascular function in other tissues. Can have both pro- and anti- inflammatory properties depending on the modality of enzymatic cleavage by different classes of proteases. Acts as a chemotactic factor for leukocyte populations expressing CMKLR1, particularly immature plasmacytoid dendritic cells, but also immature myeloid DCs, macrophages and natural killer cells. Exerts an anti-inflammatory role by preventing TNF/TNFA-induced VCAM1 expression and monocytes adhesion in vascular endothelial cells. The effect is mediated via inhibiting activation of NF-kappa-B and CRK/p38 through stimulation of AKT1/NOS3 signaling and nitric oxide production. Its dual role in inflammation and metabolism might provide a link between chronic inflammation and obesity, as well as obesity-related disorders such as type 2 diabetes and cardiovascular disease. Exhibits an antimicrobial function in the skin.

**Cellular Location** 

Secreted {ECO:0000250|UniProtKB:Q9DD06}.

#### **Tissue Location**

Expressed at the highest levels in placenta, liver, and white adipose tissue (WAT), and to a lesser extent in many other tissues such as lung, brown adipose tissue, heart, ovary, kidney, skeletal muscle and pancreas. Within WAT, expression is enriched in adipocytes as compared to the stromal vascular fraction. Expression and secretion increases dramatically with adipogenesis. Highly expressed in skin (basal and suprabasal layers of the epidermis, hair follicles and endothelial cells). Expression is elevated in numerous metabolic and inflammatory diseases including psoriasis, obesity, type 2 diabetes, metabolic syndrome and cardiovascular disease

## TIG2 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- <u>Dot Blot</u>
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>
- TIG2 Antibody Images

### TIG2 Antibody - Background

Adipocyte-secreted protein (adipokine) that regulates adipogenesis, metabolism and inflammation through activation of the chemokine-like receptor 1 (CMKLR1). Its other ligands include G protein-coupled receptor 1 (GPR1) and chemokine receptor-like 2 (CCRL2). Positively regulates adipocyte differentiation, modulates the expression of adipocyte genes involved in lipid and glucose metabolism and might play a role in angiogenesis, a process essential for the expansion of white adipose tissue. Also acts as a proinflammatory adipokine, causing an increase in secretion of proinflammatory and prodiabetic adipokines, which further impair adipose tissue metabolic function and have negative systemic effects including impaired insulin sensitivity, altered glucose and lipid metabolism, and a decrease in vascular function in other tissues. Can have both pro- and anti-inflammatory properties depending on the modality of enzymatic cleavage by different classes of proteases. Acts as a chemotactic factor for leukocyte populations expressing CMKLR1, particularly immature plasmacytoid dendritic cells, but also immature myeloid DCs, macrophages and natural killer cells. Exerts an anti-inflammatory role by preventing TNF/TNFA-induced VCAM1 expression and monocytes adhesion in vascular endothelial cells. The effect is mediated via inhibiting activation of NF-kappa-B and CRK/p38 through stimulation of AKT1/NOS3 signaling and nitric oxide production. Its dual role in inflammation and metabolism might provide a link between



chronic inflammation and obesity, as well as obesity- related disorders such as type 2 diabetes and cardiovascular disease. Exhibits an antimicrobial function in the skin.

## **TIG2 Antibody - References**

Nagpal S.,et al.J. Invest. Dermatol. 109:91-95(1997). Yokoyama-Kobayashi M.,et al.Gene 228:161-167(1999). Ota T.,et al.Nat. Genet. 36:40-45(2004). Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases. Hillier L.W.,et al.Nature 424:157-164(2003).