

# CD203c Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP52015

### **Specification**

# **CD203c Antibody - Product Information**

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW

WB, E
O14638
Human, Mouse, Rat
Rabbit
Polyclonal
100 KDa

# **CD203c Antibody - Additional Information**

#### **Gene ID 5169**

#### **Other Names**

Ectonucleotide pyrophosphatase/phosphodiesterase family member 3, E-NPP 3, Phosphodiesterase I beta, PD-Ibeta, Phosphodiesterase I/nucleotide pyrophosphatase 3, CD203c, Alkaline phosphodiesterase I, Nucleotide pyrophosphatase, NPPase, ENPP3, PDNP3

#### **Format**

0.01M PBS, pH 7.2, 0.09% (W/V) Sodium azide, Glycerol 50%

#### Storage

Store at -20 °C. Stable for 12 months from date of receipt

#### **CD203c Antibody - Protein Information**

### Name ENPP3

**Synonyms** PDNP3 {ECO:0000303|PubMed:9344668}

#### **Function**

Hydrolase that metabolizes extracellular nucleotides, including ATP, GTP, UTP and CTP (PubMed:<a href="http://www.uniprot.org/citations/29717535" target="\_blank">29717535</a>). Limits mast cell and basophil responses during inflammation and during the chronic phases of allergic responses by eliminating the extracellular ATP that functions as signaling molecule and activates basophils and mast cells and induces the release of inflammatory cytokines. Metabolizes extracellular ATP in the lumen of the small intestine, and thereby prevents ATP-induced apoptosis of intestinal plasmacytoid dendritic cells (By similarity). Has also alkaline phosphodiesterase activity (PubMed:<a href="http://www.uniprot.org/citations/11342463" target="\_blank">11342463</a>).

#### **Cellular Location**

Cell membrane; Single-pass type II membrane protein. Apical cell membrane; Single-pass type II membrane protein. Secreted Note=Detected at the cell surface of basophils (PubMed:11342463)



Detected at the apical plasma membrane of bile duct cells (PubMed:15072822). Located to the apical surface in intestinal and kidney epithelial cells. Secreted in serum, and in lumen of epithelial cells.

#### **Tissue Location**

Detected on bile ducts in liver, and in blood serum (at protein level) (PubMed:15072822). Detected in prostate and uterus (PubMed:9344668). Detected on basophils, but not neutrophils (PubMed:11342463).

### CD203c Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

### CD203c Antibody - Images

# CD203c Antibody - Background

Cleaves a variety of phosphodiester and phosphosulfate bonds including deoxynucleotides, nucleotide sugars, and NAD (By similarity).

# **CD203c Antibody - References**

Piao J.-H.,et al.Genomics 45:412-415(1997). Bechtel S.,et al.BMC Genomics 8:399-399(2007). Mungall A.J.,et al.Nature 425:805-811(2003). Mural R.J.,et al.Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases. Buehring H.J.,et al.Blood 97:3303-3305(2001).