

Rabbit Anti-WNT2 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP52104**Specification**

Rabbit Anti-WNT2 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P09544
Reactivity	Human, Mouse, Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	40418

Rabbit Anti-WNT2 Polyclonal Antibody - Additional Information**Gene ID** 7472**Other Names**

IRP; INT1L1; Protein Wnt-2; Int-1-like protein 1; Int-1-related protein; WNT2

Dilution

WB~1:100~1:500
IHC-P~1:100~1:500

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Rabbit Anti-WNT2 Polyclonal Antibody - Protein Information**Name** WNT2**Synonyms** INT1L1, IRP**Function**

Ligand for members of the frizzled family of seven transmembrane receptors. Functions in the canonical Wnt signaling pathway that results in activation of transcription factors of the TCF/LEF family (PubMed:<http://www.uniprot.org/citations/20018874> target="_blank">20018874). Functions as a upstream regulator of FGF10 expression. Plays an important role in embryonic lung development. May contribute to embryonic brain development by regulating the proliferation of dopaminergic precursors and neurons (By similarity).

Cellular Location

Secreted, extracellular space, extracellular matrix. Secreted


Tissue Location

Expressed in brain in the thalamus, in fetal and adult lung and in placenta.


Rabbit Anti-WNT2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Rabbit Anti-WNT2 Polyclonal Antibody - Images

Mouse brain lysates probed with Anti-WNT2 Polyclonal Antibody, Unconjugated (AP52104) at 1:300 in 4°C. Followed by conjugation to secondary antibody at 1:5000 90min in 37°C.



Formalin-fixed and paraffin embedded human placenta labeled with Anti-WNT2 Polyclonal Antibody, Unconjugated (AP52104) at 1:200 followed by conjugation to the secondary antibody and DAB staining

Rabbit Anti-WNT2 Polyclonal Antibody - Background

Ligand for members of the frizzled family of seven transmembrane receptors. Probable developmental protein. May be a signaling molecule which affects the development of discrete regions of tissues. Is likely to signal over only few cell diameters.