

Rabbit Anti-TIMP-2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP52136

Specification

Rabbit Anti-TIMP-2 Polyclonal Antibody - Product Information

Application WB, IHC-P Primary Accession P16035

Reactivity Human, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Calculated MW 24399

Rabbit Anti-TIMP-2 Polyclonal Antibody - Additional Information

Gene ID 7077

Other Names

DDC8; CSC-21K; KIAA1731NL; Metalloproteinase inhibitor 2; Tissue inhibitor of metalloproteinases 2; TIMP-2; TIMP2

Dilution

WB~~1:100~1:500<br \> IHC-P~~1:100~1:500

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Rabbit Anti-TIMP-2 Polyclonal Antibody - Protein Information

Name TIMP2

Function

Complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them by binding to their catalytic zinc cofactor. Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-10, MMP-13, MMP-14, MMP-15, MMP-16 and MMP-19.

Cellular Location

Secreted.


Rabbit Anti-TIMP-2 Polyclonal Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Rabbit Anti-TIMP-2 Polyclonal Antibody - Images



L1 and L2 rat lung lysates probed with Anti TIMP-2 Polyclonal Antibody, Unconjugated (AP52136) at 1:200 in 4°C. Followed by conjugation to secondary antibody at 1:3000 90min in 37°C. Predicted band 24kD. Observed band size: 21kD



Formalin-fixed and paraffin embedded rat kidney tissue labeled with Anti-TIMP-2 Polyclonal Antibody (AP52136), Unconjugated at 1:200, followed by conjugation to the secondary antibody and DAB staining

Rabbit Anti-TIMP-2 Polyclonal Antibody - Background

Complexes with metalloproteinases (such as collagenases) and irreversibly inactivates them by binding to their catalytic zinc cofactor. Known to act on MMP-1, MMP-2, MMP-3, MMP-7, MMP-8, MMP-9, MMP-13, MMP-14, MMP-15, MMP-16 and MMP-19.