

Rabbit Anti-N-cadherin Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP52149**Specification**

Rabbit Anti-N-cadherin Polyclonal Antibody - Product Information

Application	IF, WB, IHC-P
Primary Accession	P19022
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Rabbit Anti-N-cadherin Polyclonal Antibody - Additional Information**Gene ID** 1000**Other Names**

CDHN; NCAD; CD325; CDw325; Cadherin-2; Neural cadherin; N-cadherin; CDH2

Dilution

IF~1:100~1:200<br \>WB~1:100~1:500<br \>IHC-P~1:100~1:500

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

Rabbit Anti-N-cadherin Polyclonal Antibody - Protein Information**Name** CDH2**Synonyms** CDHN, NCAD**Function**

Calcium-dependent cell adhesion protein; preferentially mediates homotypic cell-cell adhesion by dimerization with a CDH2 chain from another cell. Cadherins may thus contribute to the sorting of heterogeneous cell types. Acts as a regulator of neural stem cells quiescence by mediating anchorage of neural stem cells to ependymocytes in the adult subependymal zone: upon cleavage by MMP24, CDH2-mediated anchorage is affected, leading to modulate neural stem cell quiescence. Plays a role in cell-to-cell junction formation between pancreatic beta cells and neural crest stem (NCS) cells, promoting the formation of processes by NCS cells (By similarity). Required for proper neurite branching. Required for pre- and postsynaptic organization (By similarity). CDH2 may be involved in neuronal recognition mechanism. In hippocampal neurons, may regulate dendritic spine density.


Cellular Location

Cell membrane {ECO:0000250|UniProtKB:P15116}; Single-pass type I membrane protein. Cell membrane, sarcolemma {ECO:0000250|UniProtKB:P15116}. Cell junction. Cell surface {ECO:0000250|UniProtKB:P15116}. Cell junction, desmosome {ECO:0000250|UniProtKB:P15116}. Cell junction, adherens junction {ECO:0000250|UniProtKB:P15116}. Note=Colocalizes with TMEM65 at the intercalated disk in cardiomyocytes. Colocalizes with OBSCN at the intercalated disk and at sarcolemma in cardiomyocytes {ECO:0000250|UniProtKB:P15116}


Rabbit Anti-N-cadherin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Rabbit Anti-N-cadherin Polyclonal Antibody - Images


Formalin-fixed and paraffin-embedded rat brain labeled with Anti-CDH2/N-cadherin Polyclonal Antibody, Unconjugated 1:200, overnight at 4°C, The secondary antibody was Goat Anti-Rabbit IgG, PE conjugated secondary antibody used at 1:200 dilution for 40 minutes at 37°C.



Mouse breast cancer lysates probed with Anti-N-cadherin Polyclonal Antibody, Unconjugated (AP52149) at 1:300 in 4°C. Followed by conjugation to secondary antibody at 1:5000 90min in 37°C.



Paraformaldehyde-fixed, paraffin embedded human lung carcinoma tissue; Antigen retrieval by boiling in sodium citrate buffer(pH6) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 30 minutes; Blocking buffer (normal goat serum) at 37°C for 20min; Antibody incubation with Rabbit Anti-N-cadherin Polyclonal Antibody, Unconjugated (AP52149) at 1:400 overnight at 4°C, followed by a conjugated secondary and DAB staining



Formalin-fixed and paraffin embedded rat brain labeled with Anti-CDH2/N-cadherin Polyclonal Antibody, Unconjugated at 1:200 followed by conjugation to the secondary antibody and DAB staining.

Rabbit Anti-N-cadherin Polyclonal Antibody - Background

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. Acts as a regulator of neural stem cells quiescence by mediating anchorage of neural stem cells to ependymocytes in the adult subependymal zone: upon cleavage by MMP24, CDH2-mediated anchorage is affected, leading to modulate neural stem cell quiescence. CDH2 may be involved in neuronal recognition mechanism. In hippocampal neurons, may regulate dendritic spine density (By similarity).

Rabbit Anti-N-cadherin Polyclonal Antibody - Citations

- [Exosomal circPABPC1 promotes colorectal cancer liver metastases by regulating HMGA2 in the nucleus and BMP4/ADAM19 in the cytoplasm](#)
- [Antioxidation and Antiapoptosis Characteristics of Heme Oxygenase-1 Enhance Tumorigenesis of Human Prostate Carcinoma Cells](#)
- [Inhibition of ATM reverses EMT and decreases metastatic potential of cisplatin-resistant lung cancer cells through JAK/STAT3/PD-L1 pathway.](#)
- [MiR-5100 targets TOB2 to drive epithelial-mesenchymal transition associated with activating smad2/3 in lung epithelial cells.](#)