

VE-Cadherin (Phospho-Tyr731) Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP52553

Specification

VE-Cadherin (Phospho-Tyr731) Antibody - Product Information

Application WB, IHC **Primary Accession** P33151 Reactivity Human, Mouse Host **Rabbit**

Clonality **Polyclonal** Calculated MW 87528

VE-Cadherin (Phospho-Tyr731) Antibody - Additional Information

Gene ID 1003

Other Names

Cadherin-5, 7B4 antigen, Vascular endothelial cadherin, VE-cadherin, CD144, CDH5

Dilution

WB~~1:1000 IHC~~1:50~100

Format

Rabbit IgG in phosphate buffered saline (without Mg2+ and Ca2+), pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol.

Storage Conditions

-20°C

VE-Cadherin (Phospho-Tyr731) Antibody - Protein Information

Name CDH5

Function

Cadherins are calcium-dependent cell adhesion proteins (By similarity). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types (PubMed:21269602). This

cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions (By similarity). It associates with alpha-catenin forming a link to the cytoskeleton (PubMed: 10861224). Acts in concert with KRIT1 and PALS1 to establish and maintain correct endothelial cell polarity and vascular lumen (By similarity). These effects are mediated by

recruitment and activation of the Par polarity complex and RAP1B (PubMed: 20332120). Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to



the cell junction (PubMed:20332120).

Cellular Location

Cell junction. Cell membrane; Single-pass type I membrane protein. Note=Found at cell-cell boundaries and probably at cell-matrix boundaries. KRIT1 and CDH5 reciprocally regulate their localization to endothelial cell-cell junctions.

Tissue Location


Endothelial tissues and brain.

VE-Cadherin (Phospho-Tyr731) Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

VE-Cadherin (Phospho-Tyr731) Antibody - Images



Western blot analysis of extracts from HepG2 cells, treated with Na3VO4 (0.3mM, 40mins), using VE-Cadherin (Phospho-Tyr731) antibody.





Immunohistochemistry analysis of paraffin-embedded human brain tissue using VE-Cadherin (Phospho-Tyr731) antibody.

VE-Cadherin (Phospho-Tyr731) Antibody - Background

Cadherins are calcium-dependent cell adhesion proteins. They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. This cadherin may play a important role in endothelial cell biology through control of the cohesion and organization of the intercellular junctions. It associates with alpha-catenin forming a link to the cytoskeleton. Acts in concert with KRIT1 to establish and maintain correct endothelial cell polarity and vascular lumen. These effects are mediated by recruitment and activation of the Par polarity complex and RAP1B. Required for activation of PRKCZ and for the localization of phosphorylated PRKCZ, PARD3, TIAM1 and RAP1B to the cell junction.

VE-Cadherin (Phospho-Tyr731) Antibody - References

Breviario F., et al.Arterioscler. Thromb. Vasc. Biol. 15:1229-1239(1995). Ali J., et al.Microcirculation 4:267-277(1997). Shimoyama Y., et al.Biochem. J. 349:159-167(2000). Suzuki S., et al.Cell Regul. 2:261-270(1991). Lampugnani M.G., et al.J. Cell Biol. 118:1511-1522(1992).