

**PP1C Antibody**  
**Purified Mouse Monoclonal Antibody (Mab)**  
**Catalog # AP52703****Specification**

---

**PP1C Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB                     |
| Primary Accession | <a href="#">P36873</a> |
| Reactivity        | Human                  |
| Host              | Mouse                  |
| Clonality         | Monoclonal             |
| Isotype           | IgG2b                  |
| Calculated MW     | 38 KDa                 |

**PP1C Antibody - Additional Information****Gene ID** 5501**Other Names**

EC 3.1.3.16;PP 1G;PP-1G;PP1G;PP1G\_HUMAN;PP1gamma;PPP 1G;PPP1CC;PPP1CC protein;PPP1G; Protein phosphatase 1 catalytic subunit gamma isoform;Protein phosphatase 1C catalytic subunit;Protein phosphatase 1C subunit;Protein phosphatase 2C gamma isoform; Serine/threonine phosphatase 1 gamma;Serine/threonine protein phosphatase PP1 gamma catalytic subunit;Serine/threonine-protein phosphatase PP1-gamma catalytic subunit.

**Dilution**

WB~~1:500

**Format**

Purified mouse monoclonal in buffer containing 0.1M Tris-Glycine(pH 7.4,150 mM NaCl)with 0.09% (W/V) sodium azide,0.1mg/mlBSA and 50% glycerol.

**Storage**

Store at -20 °C.Stable for 12 months from date of receipt

**PP1C Antibody - Protein Information****Name** PPP1CC**Function**

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Dephosphorylates RPS6KB1. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density- associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase.

In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E. Dephosphorylates the 'Ser-418' residue of FOXP3 in regulatory T-cells (Treg) from patients with rheumatoid arthritis, thereby inactivating FOXP3 and rendering Treg cells functionally defective (PubMed:<a href="http://www.uniprot.org/citations/23396208" target="\_blank">23396208</a>).

#### **Cellular Location**

Cytoplasm. Nucleus. Nucleus, nucleolus. Nucleus, nucleoplasm. Nucleus speckle. Chromosome, centromere, kinetochore. Cleavage furrow. Midbody Mitochondrion. Cytoplasm, cytoskeleton, microtubule organizing center Note=Colocalizes with SPZ1 in the nucleus (By similarity). Colocalizes with URI1 at mitochondrion (PubMed:17936702). Rapidly exchanges between the nucleolar, nucleoplasmic and cytoplasmic compartments (PubMed:11739654). Highly mobile in cells and can be relocalized through interaction with targeting subunits (PubMed:17965019). In the presence of PPP1R8 relocalizes from the nucleolus to nuclear speckles (PubMed:11739654). Shows a dynamic targeting to specific sites throughout the cell cycle (PubMed:12529430). Highly concentrated in nucleoli of interphase cells and localizes at kinetochores early in mitosis (PubMed:12529430). Relocalization to chromosome-containing regions occurs at the transition from early to late anaphase (PubMed:12529430). Also accumulates at the cleavage furrow and midbody by telophase (PubMed:12529430). Colocalizes with DYNLT4 in the microtubule organizing center (MTOC)(PubMed:23789093) {ECO:0000250|UniProtKB:P63087, ECO:0000269|PubMed:11739654, ECO:0000269|PubMed:12529430, ECO:0000269|PubMed:17936702, ECO:0000269|PubMed:17965019, ECO:0000269|PubMed:23789093}


#### **PP1C Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### **PP1C Antibody - Images**





Western blot detection of PP1C in HeLa, 293T and Jurkat cell lysates using PP1C mouse mAb (1:500 diluted). Predicted band size: 38KDa. Observed band size: 38KDa.

### PP1C Antibody - Background

Protein phosphatase that associates with over 200 regulatory proteins to form highly specific holoenzymes which dephosphorylate hundreds of biological targets. Protein phosphatase 1 (PP1) is essential for cell division, and participates in the regulation of glycogen metabolism, muscle contractility and protein synthesis. Dephosphorylates RPS6KB1. Involved in regulation of ionic conductances and long-term synaptic plasticity. May play an important role in dephosphorylating substrates such as the postsynaptic density- associated Ca(2+)/calmodulin dependent protein kinase II. Component of the PTW/PP1 phosphatase complex, which plays a role in the control of chromatin structure and cell cycle progression during the transition from mitosis into interphase. In balance with CSNK1D and CSNK1E, determines the circadian period length, through the regulation of the speed and rhythmicity of PER1 and PER2 phosphorylation. May dephosphorylate CSNK1D and CSNK1E.

### PP1C Antibody - References

Barker H.M., et al. Biochim. Biophys. Acta 1178:228-233(1993).  
 Bienvenut W.V., et al. Submitted (MAR-2009) to UniProtKB.  
 Bienvenut W.V., et al. Submitted (JAN-2010) to UniProtKB.  
 Norman S.A., et al. Mamm. Genome 5:41-45(1994).  
 MacKintosh R.W., et al. FEBS Lett. 371:236-240(1995).