

HDAC3 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52764**Specification**

HDAC3 Antibody - Product Information

Application	WB
Primary Accession	O15379
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	49 KDa

HDAC3 Antibody - Additional Information**Gene ID** 8841**Other Names**

HD 3;HD3;HDAC 3;HDAC3;HDAC3_HUMAN;Histone deacetylase 3 (HD3) (RPD3-2);histone deacetylase 3;RPD 3;RPD3 2;RPD3;RPD3-2;SMAP 45;SMAP45.

Dilution

WB~~1:1000

Format

ascites

Storage

Store at -20 °C.Stable for 12 months from date of receipt

HDAC3 Antibody - Protein Information**Name** HDAC3**Function**

Histone deacetylase that catalyzes the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates (PubMed:21030595, PubMed:21444723, PubMed:23911289, PubMed:25301942, PubMed:28167758, PubMed:28497810, PubMed:32404892, PubMed:22230954). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events (PubMed:<a

[23911289](http://www.uniprot.org/citations/23911289)). Histone deacetylases act via the formation of large multiprotein complexes, such as N-CoR repressor complex, which activate the histone deacetylase activity (PubMed:[23911289](http://www.uniprot.org/citations/23911289)), PubMed:[22230954](http://www.uniprot.org/citations/22230954)). Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys-27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression (PubMed:[23911289](http://www.uniprot.org/citations/23911289)). Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation (By similarity). Contributes, together with XBP1 isoform 1, to the activation of NFE2L2-mediated HMOX1 transcription factor gene expression in a PI(3)K/mTORC2/Akt-dependent signaling pathway leading to endothelial cell (EC) survival under disturbed flow/oxidative stress (PubMed:[25190803](http://www.uniprot.org/citations/25190803)). Regulates both the transcriptional activation and repression phases of the circadian clock in a deacetylase activity-independent manner (By similarity). During the activation phase, promotes the accumulation of ubiquitinated BMAL1 at the E-boxes and during the repression phase, blocks FBXL3-mediated CRY1/2 ubiquitination and promotes the interaction of CRY1 and BMAL1 (By similarity). The NCOR1-HDAC3 complex regulates the circadian expression of the core clock gene BMAL1 and the genes involved in lipid metabolism in the liver (By similarity). Also functions as a deacetylase for non-histone targets, such as KAT5, MEF2D, MAPK14, RARA and STAT3 (PubMed:[15653507](http://www.uniprot.org/citations/15653507)), PubMed:[21030595](http://www.uniprot.org/citations/21030595), PubMed:[21444723](http://www.uniprot.org/citations/21444723), PubMed:[25301942](http://www.uniprot.org/citations/25301942), PubMed:[28167758](http://www.uniprot.org/citations/28167758)). Serves as a corepressor of RARA, mediating its deacetylation and repression, leading to inhibition of RARE DNA element binding (PubMed:[28167758](http://www.uniprot.org/citations/28167758)). In association with RARA, plays a role in the repression of microRNA-10a and thereby in the inflammatory response (PubMed:[28167758](http://www.uniprot.org/citations/28167758)). In addition to protein deacetylase activity, also acts as a protein-lysine deacylase by recognizing other acyl groups: catalyzes removal of (2E)-butenoyl (crotonyl), lactoyl (lactyl) and 2-hydroxyisobutanoyl (2-hydroxyisobutyryl) acyl groups from lysine residues, leading to protein decrotonylation, delactylation and de-2-hydroxyisobutyrylation, respectively (PubMed:[28497810](http://www.uniprot.org/citations/28497810)), PubMed:[29192674](http://www.uniprot.org/citations/29192674), PubMed:[34608293](http://www.uniprot.org/citations/34608293), PubMed:[35044827](http://www.uniprot.org/citations/35044827)). Catalyzes decrotonylation of MAPRE1/EB1 (PubMed:[34608293](http://www.uniprot.org/citations/34608293)). Mediates delactylation NBN/NBS1, thereby inhibiting DNA double-strand breaks (DSBs) via homologous recombination (HR) (PubMed:[38961290](http://www.uniprot.org/citations/38961290)).

Cellular Location

Nucleus. Chromosome. Cytoplasm. Cytoplasm, cytosol. Note=Colocalizes with XBP1 and AKT1 in the cytoplasm (PubMed:25190803). Predominantly expressed in the nucleus in the presence of CCAR2 (PubMed:21030595)

Tissue Location


Widely expressed..

HDAC3 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

HDAC3 Antibody - Images



Western blot detection of HDAC3 in THP-1, LNCAP, MCF7, PC12, 293, Jurkat, 3T3 and HeLa cell lysates using HDAC3 mouse mAb (1:1000 diluted). Predicted band size: 49KDa. Observed band size: 49KDa.

HDAC3 Antibody - Background

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4), and some other non-histone substrates. Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. Histone deacetylases act via the formation of large multiprotein complexes. Participates in the BCL6 transcriptional repressor activity by deacetylating the H3 'Lys- 27' (H3K27) on enhancer elements, antagonizing EP300 acetyltransferase activity and repressing proximal gene expression. Probably participates in the regulation of transcription through its binding to the zinc-finger transcription factor YY1; increases YY1 repression activity. Required to repress transcription of the POU1F1 transcription factor. Acts as a molecular chaperone for shuttling phosphorylated NR2C1 to PML bodies for sumoylation (PubMed:21444723, PubMed:23911289). Contributes, together with XBP1 isoform 1, to the activation of NFE2L2-mediated HMOX1 transcription factor gene expression in a PI(3)K/mTORC2/Akt-dependent signaling pathway leading to endothelial cell (EC) survival under disturbed flow/oxidative stress (PubMed:25190803).

HDAC3 Antibody - References

Dangond F., et al. Biochem. Biophys. Res. Commun. 242:648-652(1998).
Yang W.-M., et al. J. Biol. Chem. 272:28001-28007(1997).
Emiliani S., et al. Proc. Natl. Acad. Sci. U.S.A. 95:2795-2800(1998).
Mahlknecht U., et al. Genomics 56:197-202(1999).
Mural R.J., et al. Submitted (SEP-2005) to the EMBL/GenBank/DDBJ databases.