

DR5 Antibody
Purified Mouse Monoclonal Antibody (Mab)
Catalog # AP52786**Specification**

DR5 Antibody - Product Information

Application	WB, ICC
Primary Accession	O14763
Reactivity	Human, Mouse
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	48 KDa

DR5 Antibody - Additional Information**Gene ID** 8795**Other Names**

Fas like protein;Apoptosis inducing protein TRICK2A/2B;Apoptosis inducing receptor TRAIL R2;CD 262;CD262;CD262 antigen;Cytotoxic TRAIL receptor 2;Death domain containing receptor for TRAIL/Apo 2L;Death domain containing receptor for TRAIL/Apo2L;Death receptor 5;DR 5;DR5;Fas like protein precursor;KILLER;KILLER/DR5;OTTHUMP00000123492; OTTHUMP00000123493;p53 regulated DNA damage inducible cell death receptor (killer);p53 regulated DNA damage inducible cell death receptor(killer);TNF related apoptosis inducing ligand receptor 2;TNF related apoptosis inducing ligand receptor 2;TNF-related apoptosis-inducing ligand receptor 2;TNFRSF10B;TR10B_HUMAN;TRAIL R2;TRAIL receptor 2;TRAIL-R2;TRAILR2;TRANCER;TRICK2;TRICK2A;TRICK2B;TRICKB;Tumor necrosis factor receptor like protein ZTNFR9;Tumor necrosis factor receptor like protein ZTNFR9;Tumor necrosis factor receptor superfamily member 10b;Tumor necrosis factor receptor superfamily, member 10b;ZTNFR9.

Dilution

WB~~1:500-1:2000

ICC~~1:100

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

DR5 Antibody - Protein Information**Name** TNFRSF10B**Synonyms** DR5, KILLER, TRAILR2, TRICK2, ZTNFR9

Function

Receptor for the cytotoxic ligand TNFSF10/TRAIL (PubMed:10549288). The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.

Cellular Location

Membrane; Single-pass type I membrane protein.


Tissue Location

Widely expressed in adult and fetal tissues; very highly expressed in tumor cell lines such as HeLaS3, K-562, HL-60, SW480, A-549 and G-361; highly expressed in heart, peripheral blood lymphocytes, liver, pancreas, spleen, thymus, prostate, ovary, uterus, placenta, testis, esophagus, stomach and throughout the intestinal tract; not detectable in brain


DR5 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DR5 Antibody - Images

Western blot detection of DR5 in Mouse kidney, Mouse brain, HCT116 and HeLa cell lysates using DR5 mouse mAb (1:500-1:2000 diluted). Predicted band size: 40/48KDa. Observed band size: 40/48KDa.



Immunocytochemistry of HeLa cells fixed by Paraformaldehyde and using DR5 mouse mAb diluted 1:100.

DR5 Antibody - Background

Receptor for the cytotoxic ligand TNFSF10/TRAIL. The adapter molecule FADD recruits caspase-8 to the activated receptor. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis. Promotes the activation of NF-kappa-B. Essential for ER stress-induced apoptosis.

DR5 Antibody - References

Screaton G.R.,et al.Curr. Biol. 7:693-696(1997).
Walczak H.,et al.EMBO J. 16:5386-5397(1997).
Schneider P.,et al.FEBS Lett. 416:329-334(1997).
Chaudhary P.M.,et al.Immunity 7:821-830(1997).
MacFarlane M.,et al.J. Biol. Chem. 272:25417-25420(1997).