

**MYLIP Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP5315c****Specification**

---

**MYLIP Antibody (Center) - Product Information**

Application	WB, FC, IHC-P,E
Primary Accession	<a href="#">Q8WY64</a>
Other Accession	<a href="#">NP_037394</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	111-139

**MYLIP Antibody (Center) - Additional Information****Gene ID** 29116**Other Names**

E3 ubiquitin-protein ligase MYLIP, 632-, Inducible degrader of the LDL-receptor, Idol, Myosin regulatory light chain interacting protein, MIR, MYLIP, BZF1, IDOL

**Target/Specificity**

This MYLIP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 111-139 amino acids from the Central region of human MYLIP.

**Dilution**

WB~~1:2000  
FC~~1:10~50  
IHC-P~~1:50~100  
E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

MYLIP Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**MYLIP Antibody (Center) - Protein Information****Name** MYLIP

**Synonyms** BZF1, IDOL

**Function** E3 ubiquitin-protein ligase that mediates ubiquitination and subsequent proteasomal degradation of myosin regulatory light chain (MRLC), LDLR, VLDLR and LRP8. Activity depends on E2 enzymes of the UBE2D family. Proteasomal degradation of MRLC leads to inhibit neurite outgrowth in presence of NGF by counteracting the stabilization of MRLC by saposin-like protein (CNPY2/MSAP) and reducing CNPY2-stimulated neurite outgrowth. Acts as a sterol-dependent inhibitor of cellular cholesterol uptake by mediating ubiquitination and subsequent degradation of LDLR.

**Cellular Location**

Cytoplasm. Cell membrane; Peripheral membrane protein


**Tissue Location**

Ubiquitously expressed.


**MYLIP Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**MYLIP Antibody (Center) - Images**


MYLIP Antibody (Center T126) (Cat. #AP5315c) western blot analysis in mouse heart tissue lysates (35ug/lane). This demonstrates the MYLIP antibody detected the MYLIP protein (arrow).




Western blot analysis of lysate from mouse kidney tissue lysate, using MYLIP Antibody (Center)(Cat. #AP5315c). AP5315c was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:10000 dilution was used as the secondary antibody. Lysate at 20ug.




All lanes : Anti-MYLIP Antibody (Center) at 1:2000 dilution Lane 1: 293 whole cell lysate Lane 2: MCF-7 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



Anti-MYLIP Antibody (Center) at 1:2000 dilution + Rat brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



MYLIP antibody (Center) (Cat. #AP5315c) immunohistochemistry analysis in formalin fixed and paraffin embedded human brain tissue followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the MYLIP antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



MYLIP Antibody (Center) (Cat. #AP5315c) flow cytometric analysis of MCF-7 cells (right histogram) compared to a negative control cell (left histogram). FITC-conjugated goat-anti-rabbit

secondary antibodies were used for the analysis.

#### **MYLIP Antibody (Center) - Background**

The ERM protein family members ezrin, radixin, and moesin are cytoskeletal effector proteins linking actin to membrane-bound proteins at the cell surface. Myosin regulatory light chain interacting protein (MYLIP) is a novel ERM-like protein that interacts with myosin regulatory light chain and inhibits neurite outgrowth.

#### **MYLIP Antibody (Center) - References**

Chasman, D.I., et al. PLoS Genet. 5 (11), E1000730 (2009)  
Lindholm, D., et al. Cell. Mol. Life Sci. 66(21):3399-3402(2009)  
Zelcer, N., et al. Science 325(5936):100-104(2009)