

CD58 Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP53294**Specification**

CD58 Antibody - Product Information

Application	WB
Primary Accession	P19256
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Calculated MW	28 KDa
Antigen Region	32-81

CD58 Antibody - Additional Information**Gene ID** 965**Other Names**

Lymphocyte function-associated antigen 3, Ag3, Surface glycoprotein LFA-3, CD58, CD58, LFA3

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human CD58. The exact sequence is proprietary.

Dilution

WB~~ 1:1000

Format

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.09% (W/V) sodium azide and 50% glycerol

Storage

Store at -20 °C.Stable for 12 months from date of receipt

CD58 Antibody - Protein Information**Name** CD58**Synonyms** LFA3**Function**

Ligand of the T-lymphocyte CD2 glycoprotein. This interaction is important in mediating thymocyte interactions with thymic epithelial cells, antigen-independent and -dependent interactions of T-lymphocytes with target cells and antigen-presenting cells and the T-lymphocyte rosetting with erythrocytes. In addition, the LFA-3/CD2 interaction may prime response by both the CD2+ and LFA-3+ cells.


Cellular Location

[Isoform 1]: Cell membrane; Single-pass type I membrane protein

CD58 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CD58 Antibody - Images

Anti-CD58 Antibody at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 28 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

CD58 Antibody - Background

Ligand of the T-lymphocyte CD2 glycoprotein. This interaction is important in mediating thymocyte interactions with thymic epithelial cells, antigen-independent and -dependent interactions of T-lymphocytes with target cells and antigen- presenting cells and the T-lymphocyte rosetting with erythrocytes. In addition, the LFA-3/CD2 interaction may prime response by both the CD2+ and LFA-3+ cells.

CD58 Antibody - References

Wallner B.P.,et al.J. Exp. Med. 166:923-932(1987).
Seed B.,et al.Nature 329:840-842(1987).
Wallich R.,et al.J. Immunol. 160:2862-2871(1998).
Ota T.,et al.Nat. Genet. 36:40-45(2004).
Ebert L.,et al.Submitted (JUN-2004) to the EMBL/GenBank/DDBJ databases.

