

Anti-Jak1 Antibody
Mouse Anti Human Monoclonal Antibody
Catalog # AP53400**Specification**

Anti-Jak1 Antibody - Product Information

Application	WB, IP
Primary Accession	P23458
Other Accession	NM_002227
Reactivity	Human, Rat
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Immunogen	Purified recombinant human Jak1 protein fragments expressed in E.coli.
Purification	Affinity purified
Calculated MW	130 KDa

Anti-Jak1 Antibody - Additional Information**Gene ID** 3716**Other Names**

JAK 1; JAK 1A; JAK 1B; JAK-1; JAK1; JAK1_HUMAN; JAK1A; JAK1B; Janus kinase 1 (a protein tyrosine kinase); Janus kinase 1; JTK3; Tyrosine protein kinase JAK 1; Tyrosine protein kinase JAK1; Tyrosine-protein kinase JAK1.

Dilution

WB~~1:500

IP~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

Anti-Jak1 Antibody - Protein Information**Name** JAK1**Synonyms** JAK1A, JAK1B**Function**

Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway (PubMed: [16239216](http://www.uniprot.org/citations/16239216)), PubMed: [28111307](http://www.uniprot.org/citations/28111307), PubMed: [32750333](http://www.uniprot.org/citations/32750333)),

PubMed:7615558, PubMed:8232552). Kinase partner for the interleukin (IL)-2 receptor (PubMed:11909529) as well as interleukin (IL)-10 receptor (PubMed:12133952). Kinase partner for the type I interferon receptor IFNAR2 (PubMed:16239216, PubMed:28111307, PubMed:32750333, PubMed:7615558, PubMed:8232552). In response to interferon-binding to IFNAR1-IFNAR2 heterodimer, phosphorylates and activates its binding partner IFNAR2, creating docking sites for STAT proteins (PubMed:7759950). Directly phosphorylates STAT proteins but also activates STAT signaling through the transactivation of other JAK kinases associated with signaling receptors (PubMed:16239216, PubMed:32750333, PubMed:8232552).

Cellular Location

Endomembrane system; Peripheral membrane protein. Note=Wholly intracellular, possibly membrane associated

Tissue Location

Expressed at higher levels in primary colon tumors than in normal colon tissue. The expression level in metastatic colon tumors is comparable to the expression level in normal colon tissue


Anti-Jak1 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Jak1 Antibody - Images





Immunoprecipitation analysis of HeLa cell lysates using Jak1 mouse mAb. Jak1 mouse mAb was used for the western blot analysis (1:500 diluted)



Western blot analysis of extracts from PC12, C6, Ramos and Jurkat cell lysates using Jak1 mouse mAb (1:500 diluted). Predicted band size: 130kDa. Observed band size: 130kDa.

Anti-Jak1 Antibody - Background

Tyrosine kinase of the non-receptor type, involved in the IFN-alpha/beta/gamma signal pathway. Kinase partner for the interleukin (IL)-2 receptor.