

### **Anti-eIF4E Antibody**

Mouse Monoclonal Antibody Catalog # AP53474

### **Specification**

## **Anti-eIF4E Antibody - Product Information**

Application WB
Primary Accession P06730
Other Accession NM 001968

Reactivity Human, Mouse, Rat

Host Mouse Clonality Monoclonal Isotype IgG1

Immunogen Purified recombinant human eif4e protein

expressed in E.coli.

Purification Affinity purified Calculated MW 25kDa KDa

## **Anti-eIF4E Antibody - Additional Information**

**Gene ID 1977** 

## **Other Names**

AUTS19; CBP; eIF 4E; eIF 4F 25 kDa subunit; EIF 4F; eIF-4E; eIF-4F 25 kDa subunit; eIF4E; EIF4E1; EIF4EL1; EIF4F; Eukaryotic translation initiation factor 4 E; Eukaryotic translation initiation factor 4E; Eukaryotic translation initiation factor 4E like 1; IF4E\_HUMAN; Messanger RNA Cap Binding Protein eIF 4E; MGC111573; mRNA cap binding protein; mRNA cap-binding protein.

### **Dilution**

WB~~1:1000

#### **Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

#### Storage

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

# **Anti-eIF4E Antibody - Protein Information**

Name EIF4E (HGNC:3287)

Synonyms EIF4EL1, EIF4F

### **Function**

Acts in the cytoplasm to initiate and regulate protein synthesis and is required in the nucleus for export of a subset of mRNAs from the nucleus to the cytoplasm which promotes processes such as RNA capping, processing and splicing (PubMed:<a

href="http://www.uniprot.org/citations/11606200" target="\_blank">11606200</a>, PubMed:<a



href="http://www.uniprot.org/citations/22684010" target="blank">22684010</a>, PubMed:<a href="http://www.uniprot.org/citations/24335285" target="blank">24335285</a>, PubMed:<a href="http://www.uniprot.org/citations/29987188" target="\_blank">29987188</a>). Component of the protein complex eIF4F, which is involved in the recognition of the mRNA cap, ATP-dependent unwinding of 5'-terminal secondary structure and recruitment of mRNA to the ribosome (By similarity). This protein recognizes and binds the 7-methylguanosine (m7G)-containing mRNA cap during an early step in the initiation of protein synthesis and facilitates ribosome binding by inducing the unwinding of the mRNAs secondary structures (PubMed: <a href="http://www.uniprot.org/citations/16271312" target="\_blank">16271312</a>, PubMed:<a href="http://www.uniprot.org/citations/22578813" target="\_blank">22578813</a>). Together with EIF4G1, antagonizes the scanning promoted by EIF1-EIF4G1 and is required for TISU translation, a process where the TISU element recognition makes scanning unnecessary (PubMed:<a href="http://www.uniprot.org/citations/29987188" target=" blank">29987188</a>). In addition to its role in translation initiation, also acts as a regulator of translation and stability in the cytoplasm (PubMed:<a href="http://www.uniprot.org/citations/24335285" target=" blank">24335285</a>). Component of the CYFIP1-EIF4E-FMR1 complex which binds to the mRNA cap and mediates translational repression: in the complex, EIF4E mediates the binding to the mRNA cap (By similarity). Component of a multiprotein complex that sequesters and represses translation of proneurogenic factors during neurogenesis (By similarity). In P-bodies, component of a complex that mediates the storage of translationally inactive mRNAs in the cytoplasm and prevents their degradation (PubMed:<a href="http://www.uniprot.org/citations/24335285" target=" blank">24335285</a>). May play an important role in spermatogenesis through translational regulation of stage-specific mRNAs during germ cell development (By similarity). As well as its roles in translation, also involved in mRNA nucleocytoplasmic transport (By similarity). Its role in mRNA export from the nucleus to the cytoplasm relies on its ability to bind the m7G cap of RNAs and on the presence of the 50-nucleotide EIF4E sensitivity element (4ESE) in the 3'UTR of sensitive transcripts (By similarity). Interaction with the 4ESE is mediated by LRPPRC which binds simultaneously to both EIF4E and the 4ESE, thereby acting as a platform for assembly for the RNA export complex (By similarity). EIF4E-dependent mRNA export is independent of ongoing protein or RNA synthesis and is also NFX1-independent but is XPO1-dependent with LRPPRC interacting with XPO1 to form an EIF4Edependent mRNA export complex (By similarity). Alters the composition of the cytoplasmic face of the nuclear pore to promote RNA export by reducing RANBP2 expression, relocalizing nucleoporin NUP214 and increasing expression of RANBP1 and RNA export factors DDX19 and GLE1 (By similarity). Promotes the nuclear export of cyclin CCND1 mRNA (By similarity). Promotes the nuclear export of NOS2/iNOS mRNA (PubMed:<a href="http://www.uniprot.org/citations/23471078" target=" blank">23471078</a>). Promotes the nuclear export of MDM2 mRNA (PubMed: <a href="http://www.uniprot.org/citations/22684010" target=" blank">22684010</a>). Promotes the export of additional mRNAs, including others involved in the cell cycle (By similarity). In the nucleus, binds to capped splice factor-encoding mRNAs and stimulates their nuclear export to enhance splice factor production by increasing their cytoplasmic availability to the translation machinery (By similarity). May also regulate splicing through interaction with the spliceosome in an RNA and m7G cap-dependent manner (By similarity). Also binds to some pre-mRNAs and may play a role in their recruitment to the spliceosome (By similarity). Promotes steady-state capping of a subset of coding and non-coding RNAs by mediating nuclear export of capping machinery mRNAs including RNMT, RNGTT and RAMAC to enhance their translation (By similarity). Stimulates mRNA 3'-end processing by promoting the expression of several core cleavage complex factors required for mRNA cleavage and polyadenylation, and may also have a direct effect through its interaction with the CPSF3 cleavage enzyme (By similarity). Rescues cells from apoptosis by promoting activation of serine/threonine- protein kinase AKT1 through mRNA export of NBS1 which potentiates AKT1 phosphorylation and also through mRNA export of AKT1 effectors, allowing for increased production of these proteins (By similarity).

href="http://www.uniprot.org/citations/22578813" target=" blank">22578813</a>, PubMed:<a

## **Cellular Location**

Cytoplasm, P-body. Cytoplasm. Cytoplasm, Stress granule. Nucleus. Nucleus speckle. Nucleus,




nuclear body Note=Interaction with EIF4ENIF1/4E-T is required for localization to processing bodies (P-bodies) (PubMed:16157702, PubMed:24335285, PubMed:25923732). Imported in the nucleus via interaction with EIF4ENIF1/4E-T via a piggy-back mechanism (PubMed:10856257) Sequestered in the nucleus by EIF4EBP1 and EIF4EBP2 (By similarity) {ECO:0000250|UniProtKB:P63073, ECO:0000269|PubMed:10856257, ECO:0000269|PubMed:16157702, ECO:0000269|PubMed:24335285, ECO:0000269|PubMed:25923732}

## **Anti-eIF4E Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **Anti-eIF4E Antibody - Images**



Western blot detection of eIF4E in MDA-MB-468, ZR75-1, MCF7, T47D and CHO-K1 cell lysates using eIF4E mouse mAb (1:1000 diluted). Predicted band size: 25KDa. Observed band size:25KDa.







Tel: 858.875.1900 Fax: 858.875.1999

Western blot detection of eIF4E in Jurkat, Hela, MCF7, K562, C6, 3T3, PC12, HepG2, CHO-K1, Raji, Raw264.7 and Ramos cell lysates using eIF4E mouse mAb (1:2000 diluted). Predicted band size: 25KDa. Observed band size:25KDa.

## Anti-eIF4E Antibody - Background

Its translation stimulation activity is repressed by binding to the complex CYFIP1-FMR1 (By similarity). Recognizes and binds the 7-methylguanosine-containing mRNA cap during an early step in the initiation of protein synthesis and facilitates ribosome bi