

**Anti-HDAC6 Antibody**  
**Mouse Monoclonal Antibody**  
**Catalog # AP53478****Specification****Anti-HDAC6 Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q9UBN7</a>
Other Accession	<a href="#">NM_006044</a>
Reactivity	Human
Host	Mouse
Clonality	Monoclonal
Isotype	IgG2a
Immunogen	Purified recombinant human HDAC6 protein expressed in E.coli.
Purification	Affinity purified
Calculated MW	160KDa KDa

**Anti-HDAC6 Antibody - Additional Information****Gene ID** 10013**Other Names**

FLJ16239 ;HD 6 ;HD6 ;HDAC 6 ;HDAC6 ;HDAC6\_HUMAN ;Histone deacetylase 6 (HD6) ;Histone deacetylase 6 ;JM 21 ;JM21 ;KIAA0901 ;OTTHUMP00000032398 ;OTTHUMP00000197663

**Dilution**

WB~~1:1000

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide, pH 7.3.

**Storage**

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze/thaw cycles.

**Anti-HDAC6 Antibody - Protein Information****Name** HDAC6 {ECO:0000303|PubMed:10220385, ECO:0000312|HGNC:HGNC:14064}**Function**

Deacetylates a wide range of non-histone substrates (PubMed:&lt;a href="http://www.uniprot.org/citations/12024216" target="\_blank"&gt;12024216&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/18606987" target="\_blank"&gt;18606987&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/20308065" target="\_blank"&gt;20308065&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/24882211" target="\_blank"&gt;24882211&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/26246421" target="\_blank"&gt;26246421&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/30538141" target="\_blank"&gt;30538141&lt;/a&gt;, PubMed:&lt;a href="http://www.uniprot.org/citations/31857589" target="\_blank"&gt;31857589&lt;/a&gt;, PubMed:&lt;a

[30770470](http://www.uniprot.org/citations/30770470), PubMed: [38534334](http://www.uniprot.org/citations/38534334), PubMed: [39567688](http://www.uniprot.org/citations/39567688)). Plays a central role in microtubule- dependent cell motility by mediating deacetylation of tubulin (PubMed: [12024216](http://www.uniprot.org/citations/12024216), PubMed: [20308065](http://www.uniprot.org/citations/20308065), PubMed: [26246421](http://www.uniprot.org/citations/26246421)). Required for cilia disassembly via deacetylation of alpha-tubulin (PubMed: [17604723](http://www.uniprot.org/citations/17604723), PubMed: [26246421](http://www.uniprot.org/citations/26246421)). Alpha-tubulin deacetylation results in destabilization of dynamic microtubules (By similarity). Promotes deacetylation of CTTN, leading to actin polymerization, promotion of autophagosome-lysosome fusion and completion of autophagy (PubMed: [30538141](http://www.uniprot.org/citations/30538141)). Deacetylates SQSTM1 (PubMed: [31857589](http://www.uniprot.org/citations/31857589)). Deacetylates peroxiredoxins PRDX1 and PRDX2, decreasing their reducing activity (PubMed: [18606987](http://www.uniprot.org/citations/18606987)). Deacetylates antiviral protein RIGI in the presence of viral mRNAs which is required for viral RNA detection by RIGI (By similarity). Sequentially deacetylates and polyubiquitinates DNA mismatch repair protein MSH2 which leads to MSH2 degradation, reducing cellular sensitivity to DNA-damaging agents and decreasing cellular DNA mismatch repair activities (PubMed: [24882211](http://www.uniprot.org/citations/24882211)). Deacetylates DNA mismatch repair protein MLH1 which prevents recruitment of the MutL alpha complex (formed by the MLH1-PMS2 heterodimer) to the MutS alpha complex (formed by the MSH2-MSH6 heterodimer), leading to tolerance of DNA damage (PubMed: [30770470](http://www.uniprot.org/citations/30770470)). Deacetylates RHOT1/MIRO1 which blocks mitochondrial transport and mediates axon growth inhibition (By similarity). Deacetylates transcription factor SP1 which leads to increased expression of ENG, positively regulating angiogenesis (PubMed: [38534334](http://www.uniprot.org/citations/38534334)). Deacetylates KHDRBS1/SAM68 which regulates alternative splicing by inhibiting the inclusion of CD44 alternate exons (PubMed: [26080397](http://www.uniprot.org/citations/26080397)). Acts as a valine sensor by binding to valine through the primate-specific SE14 repeat region (PubMed: [39567688](http://www.uniprot.org/citations/39567688)). In valine deprivation conditions, translocates from the cytoplasm to the nucleus where it deacetylates TET2 which promotes TET2-dependent DNA demethylation, leading to DNA damage (PubMed: [39567688](http://www.uniprot.org/citations/39567688)). Promotes odontoblast differentiation following IPO7-mediated nuclear import and subsequent repression of RUNX2 expression (By similarity). In addition to its protein deacetylase activity, plays a key role in the degradation of misfolded proteins: when misfolded proteins are too abundant to be degraded by the chaperone refolding system and the ubiquitin-proteasome, mediates the transport of misfolded proteins to a cytoplasmic juxtanuclear structure called aggresome (PubMed: [17846173](http://www.uniprot.org/citations/17846173)). Probably acts as an adapter that recognizes polyubiquitinated misfolded proteins and targets them to the aggresome, facilitating their clearance by autophagy (PubMed: [17846173](http://www.uniprot.org/citations/17846173)). Involved in the MTA1-mediated epigenetic regulation of ESR1 expression in breast cancer (PubMed: [24413532](http://www.uniprot.org/citations/24413532)).

### Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Nucleus. Perikaryon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, dendrite {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, axon {ECO:0000250|UniProtKB:Q9Z2V5}. Cell projection, cilium. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, cilium basal body Note=Mainly cytoplasmic where it is associated with microtubules (PubMed:12024216). Can shuttle between the cytoplasm and the nucleus (PubMed:39567688). Retained in the cytoplasm by binding to


valine via the primate-specific SE14 repeat region while valine deprivation induces nuclear localization (PubMed:39567688). Found exclusively in the cytoplasm in proliferative cells with a fraction found in the nucleus during differentiation (By similarity). May translocate to the nucleus following DNA damage (PubMed:30770470) {ECO:0000250|UniProtKB:Q9Z2V5, ECO:0000269|PubMed:12024216, ECO:0000269|PubMed:30770470, ECO:0000269|PubMed:39567688}

## Anti-HDAC6 Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## Anti-HDAC6 Antibody - Images



Western blot detection of HDAC6 in T47D, Jurkat CHO-K1(B) and CHO-K1 transfected by HDAC6-fragment EGFP fusion protein [A] cell lysates using HDAC6 mouse mAb (1:1000 diluted).

## Anti-HDAC6 Antibody - Background

Responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progr