

SFRS16 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP5351c

Specification

SFRS16 Antibody (Center) - Product Information

Application WB, FC,E
Primary Accession Q8N2M8

Other Accession <u>Q5HZB6</u>, <u>Q8CFC7</u>, <u>A0INI5</u>, <u>NP 008987.2</u>

Reactivity Human

Predicted Bovine, Mouse, Rat

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 77162
Antigen Region 254-283

SFRS16 Antibody (Center) - Additional Information

Gene ID 11129

Other Names

CLK4-associating serine/arginine rich protein, Splicing factor, arginine/serine-rich 16, Suppressor of white-apricot homolog 2, CLASRP, SFRS16, SWAP2

Target/Specificity

This SFRS16 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 254-283 amino acids from the Central region of human SFRS16.

Dilution

WB~~1:1000 FC~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SFRS16 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

SFRS16 Antibody (Center) - Protein Information

Name CLASRP



Synonyms SFRS16, SWAP2

Function Probably functions as an alternative splicing regulator. May regulate the mRNA splicing of genes such as CLK1. May act by regulating members of the CLK kinase family (By similarity).


Cellular Location Nucleus.

SFRS16 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SFRS16 Antibody (Center) - Images



SFRS16 Antibody (Center) (Cat. #AP5351c) western blot analysis in HepG2 cell line lysates (35ug/lane). This demonstrates the SFRS16 antibody detected the SFRS16 protein (arrow).





SFRS16 Antibody (Center) (Cat. #AP5351c) flow cytometric analysis of HepG2 cells (right histogram) compared to a negative control cell (left histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

SFRS16 Antibody (Center) - Background

SFRS16 is probably functions as an alternative splicing regulator. SFRS16 may regulate the mRNA splicing of genes such as CLK1. This may act by regulating members of the CLK kinase family.

SFRS16 Antibody (Center) - References

Hosgood, H.D. III, et al. Occup Environ Med 66(12):848-853(2009) Sugiyama, N., et al. Mol. Cell Proteomics 6(6):1103-1109(2007) Olsen, J.V., et al. Cell 127(3):635-648(2006)