

Anti-HCAP (pS1083) Antibody
Catalog # AP53817**Specification**

Anti-HCAP (pS1083) Antibody - Product Information

Application	WB
Primary Accession	Q9UQE7
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	141542

Anti-HCAP (pS1083) Antibody - Additional Information**Gene ID** 9126**Other Names**

BAM; BMH; CSPG6; SMC3L1; Structural maintenance of chromosomes protein 3; SMC protein 3; SMC-3; Basement membrane-associated chondroitin proteoglycan; Bamacan; Chondroitin sulfate proteoglycan 6; Chromosome-associated polypeptide; hCAP

Target/Specificity

Recognizes endogenous levels of HCAP (pS1083) protein.

Dilution

WB~~1/500 - 1/1000

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-HCAP (pS1083) Antibody - Protein Information**Name** SMC3**Synonyms** BAM, BMH, CSPG6, SMC3L1**Function**

Central component of cohesin, a complex required for chromosome cohesion during the cell cycle. The cohesin complex may form a large proteinaceous ring within which sister chromatids can be trapped. At anaphase, the complex is cleaved and dissociates from chromatin, allowing sister chromatids to segregate. Cohesion is coupled to DNA replication and is involved in DNA repair. The cohesin complex also plays an important role in spindle pole assembly during mitosis and in chromosomes movement.

Cellular Location


Nucleus {ECO:0000250|UniProtKB:Q9CW03}. Chromosome {ECO:0000250|UniProtKB:Q9CW03}. Chromosome, centromere {ECO:0000250|UniProtKB:Q9CW03}. Note=Associates with chromatin. Before prophase it is scattered along chromosome arms. During prophase, most of cohesin complexes dissociate from chromatin probably because of phosphorylation by PLK, except at centromeres, where cohesin complexes remain. At anaphase, the RAD21 subunit of the cohesin complex is cleaved, leading to the dissociation of the complex from chromosomes, allowing chromosome separation. The phosphorylated form at Ser-1083 is preferentially associated with unsynapsed chromosomal regions (By similarity). {ECO:0000250|UniProtKB:Q9CW03}

Anti-HCAP (pS1083) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-HCAP (pS1083) Antibody - Images



Western blot analysis of HCAP (pS1083) expression in A549 (A), NS-1 (B), PC12 (C) whole cell lysates.

Anti-HCAP (pS1083) Antibody - Background

Rabbit polyclonal antibody to HCAP (pS1083)