

Anti-ZNF496 Antibody
Catalog # AP53838**Specification**

Anti-ZNF496 Antibody - Product Information

Application	WB
Primary Accession	Q96IT1
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	66908

Anti-ZNF496 Antibody - Additional Information**Gene ID** 84838**Other Names**

ZKSCAN17; Zinc finger protein 496; Zinc finger protein with KRAB and SCAN domains 17

Target/Specificity

Recognizes endogenous levels of ZNF496 protein.

Dilution

WB~~1/500 - 1/1000

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-ZNF496 Antibody - Protein Information**Name** ZNF496**Synonyms** ZKSCAN17**Function**

DNA-binding transcription factor that can both act as an activator and a repressor.

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187}.

Anti-ZNF496 Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-ZNF496 Antibody - Images



Western blot analysis of ZNF496 expression in HEK293T (A), Raw264.7 (B), PC12 (C) whole cell lysates.



Immunohistochemical analysis of ZNF496 staining in human lung formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. AEC was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-ZNF496 Antibody - Background

Rabbit polyclonal antibody to ZNF496