

Anti-TAGAP Antibody
Catalog # AP53934**Specification**

Anti-TAGAP Antibody - Product Information

Application	WB, IHC
Primary Accession	Q8N103
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	80703

Anti-TAGAP Antibody - Additional Information**Gene ID** 117289**Other Names**

TAGAP1; T-cell activation Rho GTPase-activating protein; T-cell activation GTPase-activating protein

Target/Specificity

Recognizes endogenous levels of TAGAP protein.

Dilution

WB~~1/500 - 1/2000

IHC~~1:100~500

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-TAGAP Antibody - Protein Information**Name** TAGAP**Synonyms** TAGAP1**Function**


May function as a GTPase-activating protein and may play important roles during T-cell activation.

Anti-TAGAP Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-TAGAP Antibody - Images



Western blot analysis of TAGAP expression in mouse kidney (A), rat kidney (B), rat liver (C), rat heart (D) whole cell lysates.



Immunohistochemical analysis of TAGAP staining in human breast cancer formalin fixed paraffin embedded tissue section. The section was pre-treated using heat mediated antigen retrieval with sodium citrate buffer (pH 6.0). The section was then incubated with the antibody at room temperature and detected using an HRP conjugated compact polymer system. DAB was used as the chromogen. The section was then counterstained with haematoxylin and mounted with DPX.

Anti-TAGAP Antibody - Background

Rabbit polyclonal antibody to TAGAP