

Anti-Histone H2B (AcK85) Antibody
Catalog # AP54016**Specification****Anti-Histone H2B (AcK85) Antibody - Product Information**

Application	WB
Primary Accession	P33778
Other Accession	P62807 , P58876 , Q93079 , P06899 , O60814 , Q99880 , Q99879 , Q99877 , P23527
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	13950

Anti-Histone H2B (AcK85) Antibody - Additional Information**Gene ID** 3018**Other Names**

HIST1H2BB; H2BFF; Histone H2B type 1-B; Histone H2B.1; Histone H2B.f; H2B/f; HIST1H2BC; H2BFL; HIST1H2BE; H2BFH; HIST1H2BF; H2BFG; HIST1H2BG; H2BFA; HIST1H2BI; H2BFB; Histone H2B type 1-C/E/F/G/I; Histone H2B.1 A; Histone H2B.a; H2B/a; Histone H2B.g; H2B/g; Histone H2B.h; H2B/h; Histone H2B.k; H2B/k; Histone H2B.l; H2B/l; HIST1H2BD; H2BFB; HIRIP2; Histone H2B type 1-D; HIRA-interacting protein 2; Histone H2B.1 B; Histone H2B.b; H2B/b; HIST1H2BH; H2BFJ; Histone H2B type 1-H; Histone H2B.j; H2B/j; HIST1H2BJ; H2BFR; Histone H2B type 1-J; Histone H2B.1; Histone H2B.r; H2B/r; HIST1H2BK; H2BFT; HIRIP1; Histone H2B type 1-K; H2B K; HIRA-interacting protein 1; HIST1H2BL; H2BFC; Histone H2B type 1-L; Histone H2B.c; H2B/c; HIST1H2BM; H2BFE; Histone H2B type 1-M; Histone H2B.e; H2B/e; HIST1H2BN; H2BFD; Histone H2B type 1-N; Histone H2B.d; H2B/d; HIST1H2BO; H2BFH; H2BFN; Histone H2B type 1-O; Histone H2B.2; Histone H2B.n; H2B/n

Target/Specificity

KLH-conjugated synthetic peptide encompassing a sequence within the center region of human Histone H2B (AcK85). The exact sequence is proprietary.

Dilution

WB~~1/500 - 1/1000

Format

Liquid in 0.42% Potassium phosphate, 0.87% Sodium chloride, pH 7.3, 30% glycerol, and 0.09% (W/V) sodium azide.

Storage

Store at -20 °C.Stable for 12 months from date of receipt

Anti-Histone H2B (AcK85) Antibody - Protein Information**Name** H2BC3 ([HGNC:4751](#))

Function

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.


Cellular Location

Nucleus. Chromosome.

Anti-Histone H2B (AcK85) Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Anti-Histone H2B (AcK85) Antibody - Images

Western blot analysis of Histone H2B (AcK85) expression in mouse kidney (A) whole cell lysates.

Anti-Histone H2B (AcK85) Antibody - Background

Rabbit polyclonal antibody to Histone H2B (AcK85)