



NCoR1 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54187

Specification

NCoR1 Polyclonal Antibody - Product Information

Application
Primary Accession
Reactivity
Host
Clonality
Calculated MW
Physical State
Immunogen

Epitope Specificity Isotype **Purity** affinity purified by Protein A

Buffer

SUBCELLULAR LOCATION SIMILARITY

SUBUNIT

IHC-P, IHC-F, IF, E

O75376 Rat Rabbit Polyclonal 271 KDa Liquid

KLH conjugated synthetic peptide derived

from mouse NC0R1 2301-2400/2453

IqG

0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.

Nucleus

Belongs to the N-CoR nuclear receptor corepressors family. Contains 2 SANT

domains.

Interacts with C1D, SIAH2, HDAC7, SAP30, SIN3A and SIN3B (By similarity). Forms a large corepressor complex that contains. SIN3A/B and histone deacetylases HDAC1 and HDAC2. This complex associates with the thyroid (TR) and the retinoid acid receptors (RAR) in the absence of ligand. Interacts directly with RARA; the interaction is faciliated with RARA trimethylation. Interacts with DACH1. Component of the N-Cor repressor complex, at least composed of NCOR1, NCOR2, HDAC3, TBL1X, TBL1XR1, CORO2A and GPS2. Interacts with TRIM28 and KDM3A. Interacts with ZBTB33; the interaction serves to recruit the N-CoR complex to promoter regions containing methylated CpG dinucleotides. Interacts with HDAC9 (via its catalytic domain). Interacts with CBFA2T3 and HEXIM1. Interacts (via the RRFD1 domain) with **BAZ1A** (via its N-terminal); the interaction corepresses a number of NCOR1-regulated genes.

Ubiquitinated; mediated by SIAH2 and

Post-translational modifications



Important Note

leading to its subsequent proteasomal degradation.

This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Nuclear co-repressor 2 (N-CoR2) gene (NCOR2, previously called silencing mediator for retinoid and thyroid hormone receptor SMRT) is recruited to nuclear and non-nuclear receptors in a large repressing complex containing also N-CoR1, mSin3 and HDACs. This large complex represses transcription in absence of ligand. Mediates the transcriptional repression activity of some nuclear receptors by promoting chromatin condensation, thus preventing access of the basal transcription. Tissue specificity: Ubiquitous. It is belongs to the N-CoR nuclear receptor corepressors family.

NCoR1 Polyclonal Antibody - Additional Information

Gene ID 9611

Other Names

Nuclear receptor corepressor 1, N-CoR, N-CoR1, NCOR1, KIAA1047

Dilution

IHC-P~~N/A<br \> <span class
="dilution_IHC-F">IHC-F~~N/A<br \> <span class
="dilution_IF">IF~~1:50~200<br \> E~~N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

NCoR1 Polyclonal Antibody - Protein Information

Name NCOR1

Synonyms KIAA1047

Function

Mediates transcriptional repression by certain nuclear receptors (PubMed:20812024). Part of a complex which promotes histone deacetylation and the formation of repressive chromatin structures which may impede the access of basal transcription factors. Participates in the transcriptional repressor activity produced by BCL6. Recruited by ZBTB7A to the androgen response elements/ARE on target genes, negatively regulates androgen receptor signaling and androgen-induced cell proliferation (PubMed:20812024). Mediates the NR1D1-dependent repression and circadian regulation of TSHB expression (By similarity). The NCOR1-HDAC3 complex regulates the circadian expression of the core clock gene ARTNL/BMAL1 and the genes involved in lipid metabolism in the liver (By similarity).

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00624}.



Tel: 858.875.1900 Fax: 858.875.1999



NCoR1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

NCoR1 Polyclonal Antibody - Images