

MYPN Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54252

Specification

MYPN Polyclonal Antibody - Product Information

Application	IHC-P
Primary Accession	Q86TC9
Reactivity	Rat, Pig, Dog, Bovine
Host	Rabbit
Clonality	Polyclonal
Calculated MW	145257

MYPN Polyclonal Antibody - Additional Information

Gene ID 84665

Other Names

Myopalladin, 145 kDa sarcomeric protein, MYPN, MYOP

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

MYPN Polyclonal Antibody - Protein Information

Name MYPN

Synonyms MYOP

Function

Component of the sarcomere that tethers together nebulin (skeletal muscle) and nebulin (cardiac muscle) to alpha-actinin, at the Z lines.

Cellular Location

Cytoplasm. Nucleus. Cytoplasm, myofibril, sarcomere. Cytoplasm, myofibril, sarcomere, Z line.
Note=Bound to sarcomere both at the Z-line periphery and in the central I-band region.

Tissue Location


Expressed in adult skeletal muscle and fetal heart.

MYPN Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


MYPN Polyclonal Antibody - Images



Tissue/cell: Rat skeletal muscle; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-MYPN Polyclonal Antibody, Unconjugated(bs-10366R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining



Tissue/cell: Mouse embryos; 4% Paraformaldehyde-fixed and paraffin-embedded;

Antigen retrieval: citrate buffer (0.01M, pH 6.0), Boiling bathing for 15min; Block endogenous peroxidase by 3% Hydrogen peroxide for 30min; Blocking buffer (normal goat serum,C-0005) at 37°C for 20 min;

Incubation: Anti-MYPN Polyclonal Antibody, Unconjugated(bs-10366R) 1:200, overnight at 4°C, followed by conjugation to the secondary antibody(SP-0023) and DAB(C-0010) staining