

ATP6V1G2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54324

Specification

ATP6V1G2 Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW Physical State Immunogen Epitope Specificity Isotype Purity affinity purified by Protein A	WB, IHC-P, IHC-F, IF, ICC, E <u>O95670</u> Rat, Pig, Dog, Bovine Rabbit Polyclonal 14 KDa Liquid KLH conjugated synthetic peptide derived from human ATP6V1G2 51-118/118 IgG
Buffer	0.01M TBS (pH7.4) with 1% BSA, 0.02% Proclin300 and 50% Glycerol.
SUBCELLULAR LOCATION	Melanosome. Note=Highly enriched in late-stage melanosomes.
SIMILARITY	Belongs to the V-ATPase G subunit family.
SUBUNIT	V-ATPase is a heteromultimeric enzyme composed of a peripheral catalytic V1 complex (components A to H) attached to an integral membrane V0 proton pore complex (components: a, c, c', c'' and d).
Important Note	This product as supplied is intended for research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of intracellular compartments of eukaryotic cells. V-ATPase dependent acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A and three B subunits, two G subunits plus the C, D, E, F, and H subunits. The V1 domain contains the ATP catalytic site. The V0 domain consists of five different subunits: a, c, c', c'', and d. Additional isoforms of many of the V1 and V0 subunit proteins are encoded by multiple genes or alternatively spliced transcript variants. This encoded protein is one of three V1 domain G subunit proteins. This gene had previous gene symbols of ATP6G and ATP6G2. Alternatively spliced transcription also exists between this gene and the downstream DEAD (Asp-Glu-Ala-Asp) box polypeptide 39B (DDX39B) gene. [provided by RefSeq, Feb 2011]

ATP6V1G2 Polyclonal Antibody - Additional Information



Gene ID 534

Other Names V-type proton ATPase subunit G 2, V-ATPase subunit G 2, V-ATPase 13 kDa subunit 2, Vacuolar proton pump subunit G 2, ATP6V1G2, ATP6G, ATP6G2, NG38

Target/Specificity Brain.

Dilution WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-F~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ATP6V1G2 Polyclonal Antibody - Protein Information

Name ATP6V1G2

Synonyms ATP6G, ATP6G2, NG38

Function

Subunit of the V1 complex of vacuolar(H+)-ATPase (V-ATPase), a multisubunit enzyme composed of a peripheral complex (V1) that hydrolyzes ATP and a membrane integral complex (V0) that translocates protons. V-ATPase is responsible for acidifying and maintaining the pH of intracellular compartments and in some cell types, is targeted to the plasma membrane, where it is responsible for acidifying the extracellular environment.

Cellular Location

Melanosome. Cytoplasmic vesicle, clathrin-coated vesicle membrane {ECO:0000250|UniProtKB:Q0VCV6}; Peripheral membrane protein. Note=Highly enriched in late-stage melanosomes

Tissue Location Brain..

ATP6V1G2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence



Immunoprecipitation

- Flow Cytomety
- <u>Cell Culture</u>

ATP6V1G2 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (Mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ATP6V1G2) Polyclonal Antibody, Unconjugated (bs-10930R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.



Paraformaldehyde-fixed, paraffin embedded (Rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ATP6V1G2) Polyclonal Antibody, Unconjugated (bs-10930R) at 1:500 overnight at 4°C, followed by a conjugated secondary (sp-0023) for 20 minutes and DAB staining.