

MDGA2 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54370

Specification

MDGA2 Polyclonal Antibody - Product Information

Application Primary Accession

Reactivity
Host
Clonality
Calculated MW
Physical State
Immunogen

Epitope Specificity

Isotype **Purity**

affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

WB, IHC-P, IHC-F, IF, ICC, E

O7Z553

Rabbit

Liquid

laG

Polyclonal

751-850/956

102 KDa

Rat, Dog, Bovine

from human MDGA2

SUBCELLULAR LOCATION Cell membrane.

SIMILARITY Contains 6 Ig-like (immunoglobulin-like)

domains. Contains 1 MAM domain.

Important Note

This product as supplied is intended for research use only, not for use in human,

therapeutic or diagnostic applications.

KLH conjugated synthetic peptide derived

Background Descriptions

MDGA2 is a cell membrane protein which contains six Ig-like (immunoglobulin-like) domains and one MAM domain. Analyses of the full-length coding region of MDGA1 and MDGA2 indicate that they encode proteins that comprise a novel subgroup of the Ig superfamily and have a unique structural organization consisting of six immunoglobulin (Ig)-like domains followed by a single MAM domain. Biochemical characterization demonstrates that MDGA1 and MDGA2 proteins are highly glycosylated, and that MDGA1 is tethered to the cell membrane by a GPI anchor. The MDGAs are differentially expressed by subpopulations of neurons in both the central and peripheral nervous systems, including neurons of the basilar pons, inferior olive, cerebellum, cerebral cortex, olfactory bulb, spinal cord, and dorsal root and trigeminal ganglia. The similarity of MDGAs to other Ig-containing molecules and their temporal-spatial patterns of expression within restricted neuronal populations, for example migrating pontine neurons and D1 spinal interneurons, suggest a role for these novel proteins in regulating neuronal migration, as well as other aspects of neural development, including axon guidance.

MDGA2 Polyclonal Antibody - Additional Information

Gene ID 161357

Other Names

MAM domain-containing glycosylphosphatidylinositol anchor protein 2, MAM domain-containing



protein 1, MDGA2, MAMDC1

Dilution

WB~~1:1000<br \><span class
="dilution_IHC-P">IHC-P~~N/A<br \><span class
="dilution_IHC-F">IHC-F~~N/A<br \><span class
="dilution_IF">IF~~1:50~200<br \>ICC~~N/A<br \><br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A</pa></pr>

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

MDGA2 Polyclonal Antibody - Protein Information

\>E~~N/A

Name MDGA2

Synonyms MAMDC1

Function

May be involved in cell-cell interactions.

Cellular Location

Cell membrane; Lipid-anchor, GPI- anchor

Tissue Location

Detected in Leydig cells, syncytiotrophoblast, duodenal villi epithelial cells and neutrophils from kidney and cutaneous squamous cell carcinoma (at protein level)

MDGA2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MDGA2 Polyclonal Antibody - Images