

PCDHGC3 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) **Catalog # AP54407**

Specification

PCDHGC3 Polyclonal Antibody - Product Information

WB, IHC-P, IHC-F, IF, ICC, E Application

Primary Accession 09UN70 Reactivity Rat Host Rabbit Clonality **Polyclonal** Calculated MW 98 KDa

Physical State Immunogen KLH conjugated synthetic peptide derived

Liquid

laG

from human PCDHGC3/PCDH2

251-350/934 **Epitope Specificity**

Isotype **Purity** affinity purified by Protein A

Buffer 0.01M TBS (pH7.4) with 1% BSA, 0.02%

Proclin300 and 50% Glycerol.

SUBCELLULAR LOCATION Cell membrane. Single-pass type I

membrane protein.

Contains 6 cadherin domains. **SIMILARITY**

This product as supplied is intended for Important Note research use only, not for use in human, therapeutic or diagnostic applications.

Background Descriptions

Protocadherins are a large family of cadherin-like cell adhesion proteins that are involved in the establishment and maintenance of neuronal connections in the brain. There are three protocadherin gene clusters, designated alpha, beta and gamma, all of which contain multiple tandemly arranged genes. PCDH2 (protocadherin-2), also known as PCDHGC3 (protocadherin gamma subfamily C, 3) or PC43, is a 934 amino acid single-pass type I membrane protein that contains six cadherin domains and belongs to the protocadherin gamma family. Functioning as a calcium-dependent cell-adhesion protein, PCDH2 is thought to be involved in the establishment and maintenance of neuronal connections within the brain. Multiple isoforms of PCDH2 exist due to alternative splicing events.

PCDHGC3 Polyclonal Antibody - Additional Information

Gene ID 5098

Other Names

Protocadherin gamma-C3, PCDH-gamma-C3, Protocadherin-2, Protocadherin-43, PC-43, PCDHGC3, PCDH2

Dilution

WB~~1:1000<br \><span class</pre>



="dilution_IHC-P">IHC-P~~N/A<br \>IHC-F~~N/A<br \><span class

="dilution_IF">IF \sim 1:50 \sim 200<br \>ICC \sim N/A<br \>E \sim N/A

Storage

Store at -20 $^{\circ}$ C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 $^{\circ}$ C.

PCDHGC3 Polyclonal Antibody - Protein Information

Name PCDHGC3

Synonyms PCDH2

Function

Potential calcium-dependent cell-adhesion protein. May be involved in the establishment and maintenance of specific neuronal connections in the brain.

Cellular Location

Cell membrane; Single-pass type I membrane protein

PCDHGC3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

PCDHGC3 Polyclonal Antibody - Images