

SPRR3 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54409**Specification**

SPRR3 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9UBC9
Reactivity	Rat
Host	Rabbit
Clonality	Polyclonal
Calculated MW	18154

SPRR3 Polyclonal Antibody - Additional Information**Gene ID** 6707**Other Names**

Small proline-rich protein 3, 22 kDa pancornulin, Cornifin beta, Esophagin, SPRR3, SPRC

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

SPRR3 Polyclonal Antibody - Protein Information**Name** SPRR3**Synonyms** SPRC**Function**

Cross-linked envelope protein of keratinocytes.

Cellular Location

Cytoplasm.


SPRR3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)

- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SPRR3 Polyclonal Antibody - Images



Sample:


U937(Human) Cell Lysate at 30 ug

Primary: Anti- SPRR3 (bs-11163R) at 1/1000 dilution

Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution

Predicted band size: 18 kD

Observed band size: 18 kD



Paraformaldehyde-fixed, paraffin embedded (Human esophageal cancer); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (SPRR3) Polyclonal Antibody, Unconjugated (bs-11163R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.