

## EPS15 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54434

#### Specification

# **EPS15 Polyclonal Antibody - Product Information**

Application Primary Accession Reactivity Host Clonality Calculated MW IHC-P, IHC-F, IF, ICC, E P42566 Rat, Pig Rabbit Polyclonal 98656

## **EPS15 Polyclonal Antibody - Additional Information**

Gene ID 2060

Other Names Epidermal growth factor receptor substrate 15, Protein Eps15, Protein AF-1p, EPS15, AF1P

Dilution <span class ="dilution\_IHC-P">IHC-P~~N/A</span><br \><span class ="dilution\_IHC-F">IHC-F~~N/A</span><br \><span class ="dilution\_IF">IF~~1:50~200</span><br \><span class ="dilution\_ICC">ICC~~N/A</span><br \><span class ="dilution\_E">E~~N/A</span>

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

**Storage** Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

# **EPS15** Polyclonal Antibody - Protein Information

Name EPS15

Synonyms AF1P

Function

Involved in cell growth regulation. May be involved in the regulation of mitogenic signals and control of cell proliferation. Involved in the internalization of ligand-inducible receptors of the receptor tyrosine kinase (RTK) type, in particular EGFR. Plays a role in the assembly of clathrin-coated pits (CCPs). Acts as a clathrin adapter required for post-Golgi trafficking. Seems to be involved in CCPs maturation including invagination or budding. Involved in endocytosis of integrin beta-1 (ITGB1) and transferrin receptor (TFR); internalization of ITGB1 as DAB2-dependent cargo but not TFR seems to require association with DAB2.



#### **Cellular Location**

Cytoplasm. Cell membrane; Peripheral membrane protein; Cytoplasmic side. Membrane, clathrin-coated pit Note=Recruited to the plasma membrane upon EGFR activation and localizes to coated pits. Colocalizes with UBQLN1 in ubiquitin-rich cytoplasmic aggregates that are not endocytic compartments and in cytoplasmic juxtanuclear structures called aggresomes

**Tissue Location** Ubiquitously expressed.

## **EPS15 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

**EPS15 Polyclonal Antibody - Images**