

ELAVL4 Polyclonal Antibody

Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP54453

Specification

ELAVL4 Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality Calculated MW WB, IHC-P, IHC-F, IF, ICC, E <u>P26378</u> Rat, Dog, Bovine Rabbit Polyclonal 42398

ELAVL4 Polyclonal Antibody - Additional Information

Gene ID 1996

Other Names

ELAV-like protein 4, Hu-antigen D, HuD, Paraneoplastic encephalomyelitis antigen HuD, ELAVL4, HUD, PNEM

Dilution WB~~1:1000<br \>IHC-P~~N/A<br \>IHC-P~~N/A<br \>IF~~1:50~200<br \>ICC~~N/A<br \>E~~N/A<br \>ICC~~N/A<br \>ICC~~N/A<br \>ICC~~N/A</span

Format 0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

ELAVL4 Polyclonal Antibody - Protein Information

Name ELAVL4

Synonyms HUD, PNEM

Function

```
RNA-binding protein that is involved in the post- transcriptional regulation of mRNAs (PubMed:<a href="http://www.uniprot.org/citations/10710437" target="_blank">10710437</a>, PubMed:<a href="http://www.uniprot.org/citations/12034726" target="_blank">12034726</a>, PubMed:<a href="http://www.uniprot.org/citations/12468554" target="_blank">12468554</a>, PubMed:<a href="http://www.uniprot.org/citations/12468554" target="_blank">12468554</a>, PubMed:<a href="http://www.uniprot.org/citations/12468554" target="_blank">12468554</a>, PubMed:<a href="http://www.uniprot.org/citations/17035636" target="_blank">17035636</a>, PubMed:<a href="http://www.uniprot.org/citations/17035636" target="_blank">17035636</a>, PubMed:<a href="http://www.uniprot.org/citations/17035636" target="_blank">17035636</a>, PubMed:<a href="http://www.uniprot.org/citations/17234598" target="_blank">17234598</a>, P
```

href="http://www.uniprot.org/citations/7898713" target="_blank">7898713). Plays a role in the regulation of mRNA stability, alternative splicing and translation (PubMed:10710437, PubMed:12034726, PubMed:12468554, PubMed:17035636, PubMed:17035636, PubMed:17035636, PubMed:17035636, PubMed:17234598, PubMed:17234598, PubMed:1703636, PubMed:1703636, PubMed:<a href="http:

href="http://www.uniprot.org/citations/10710437" target="_blank">10710437, PubMed:12034726, PubMed:12468554, PubMed:12468554, PubMed:7898713). Many of the target mRNAs are coding for RNA-binding proteins, transcription factors and proteins involved in RNA processing and/or neuronal development and function (By similarity). By binding to the mRNA 3'UTR, decreases mRNA deadenylation and thereby contributes to the stabilization of mRNA molecules and their protection from decay (PubMed:12034726). Also binds to the polyadenylated (poly(A)) tail in the 3'UTR of mRNA, thereby increasing its affinity for mRNA binding (PubMed:<a href="http://www.uniprot.org/citations/12034726"

target=" blank">12034726). Mainly plays a role in neuron-specific RNA processing by stabilization of mRNAs such as GAP43, ACHE and mRNAs of other neuronal proteins, thereby contributing to the differentiation of neural progenitor cells, nervous system development, learning and memory mechanisms (PubMed:12034726, PubMed:12468554, PubMed:17234598, PubMed:18218628). Involved in the negative regulation of the proliferative activity of neuronal stem cells and in the positive regulation of neuronal differentiation of neural progenitor cells (By similarity). Promotes neuronal differentiation of neural stem/progenitor cells in the adult subventricular zone of the hippocampus by binding to and stabilizing SATB1 mRNA (By similarity). Binds and stabilizes MSI1 mRNA in neural stem cells (By similarity). Exhibits increased binding to ACHE mRNA during neuronal differentiation, thereby stabilizing ACHE mRNA and enhancing its expression (PubMed:12468554, PubMed:17234598). Protects CDKN1A mRNA from decay by binding to its 3'-UTR (By similarity). May bind to APP and BACE1 mRNAS and the BACE1AS IncRNA and enhance their stabilization (PubMed:24857657). Plays a role in neurite outgrowth and in the establishment and maturation of dendritic arbors, thereby contributing to neocortical and hippocampal circuitry

function (By similarity). Stabilizes GAP43 mRNA and protects it from decay during postembryonic development in the brain (PubMed:12034726). By promoting the stabilization of GAP43 mRNA, plays a role in NGF-mediated neurite outgrowth (By similarity). Binds to BDNF long 3'UTR mRNA, thereby leading to its stabilization and increased dendritic translation after activation of PKC (By similarity). By increasing translation of BDNF after nerve injury, may contribute to nerve regeneration (By similarity). Acts as a stabilizing factor by binding to the 3'UTR of NOVA1 mRNA, thereby increasing its translation and enhancing its functional activity in neuron-specific splicing (PubMed:18218628). Stimulates translation of mRNA in a poly(A)- and cap-dependent manner, possibly by associating with the EIF4F cap-binding complex (By similarity). May also negatively regulate translation by binding to the 5'UTR of Ins2 mRNA, thereby repressing its translation (By similarity). Upon glucose stimulation, Ins2 mRNA is released from ELAVL4 and translational inhibition is abolished (By similarity). Also plays a role in the regulation of alternative splicing (PubMed: 17035636). May regulate alternative splicing of CALCA pre-mRNA into Calcitonin and Calcitonin gene-related peptide 1



(CGRP) by competing with splicing regulator TIAR for binding to U-rich intronic sequences of CALCA pre- mRNA (PubMed:17035636).

Cellular Location

Cytoplasm. Perikaryon {ECO:0000250|UniProtKB:009032}. Cell projection, dendrite {ECO:0000250|UniProtKB:009032}. Cell projection, axon {ECO:0000250|UniProtKB:Q61701}. Cell projection, growth cone {ECO:0000250|UniProtKB:Q61701}. Note=Co-localizes with ribosomal RNA in polysomes. {ECO:0000250|UniProtKB:009032}

Tissue Location

Expressed in pancreatic beta cells (at protein level) (PubMed:22387028). Expressed in the brain (PubMed:14702039, PubMed:1655278).

ELAVL4 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

ELAVL4 Polyclonal Antibody - Images



Paraformaldehyde-fixed, paraffin embedded (mouse brain); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (ELAVL4) Polyclonal Antibody, Unconjugated (bs-11319R) at 1:200 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructionsand DAB staining.





Sample:

SY5Y(Human) Cell Lysate at 30 ug Primary: Anti- ELAVL4 (bs-11319R) at 1/1000 dilution Secondary: IRDye800CW Goat Anti-Rabbit IgG at 1/20000 dilution Predicted band size: 42 kD Observed band size: 42 kD