

ATP11C Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP5446c

Specification

ATP11C Antibody (Center) - Product Information

| | |
|-------------------|--------------------------------|
| Application | FC, IHC-P, WB,E |
| Primary Accession | Q8NB49 |
| Other Accession | NP_001010986.1 |
| Reactivity | Human, Hamster, Mouse |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 129477 |
| Antigen Region | 589-616 |

ATP11C Antibody (Center) - Additional Information

Gene ID 286410

Other Names

Phospholipid-transporting ATPase IG, ATPase IQ, ATPase class VI type 11C, P4-ATPase flippase complex alpha subunit ATP11C, ATP11C, ATPIG, ATPIQ

Target/Specificity

This ATP11C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 589-616 amino acids from the Central region of human ATP11C.

Dilution

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ATP11C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ATP11C Antibody (Center) - Protein Information

Name ATP11C {ECO:0000303|PubMed:26944472}

Synonyms ATPIG, ATPIQ

Function Catalytic component of a P4-ATPase flippase complex which catalyzes the hydrolysis of ATP coupled to the transport of aminophospholipids, phosphatidylserines (PS) and phosphatidylethanolamines (PE), from the outer to the inner leaflet of the plasma membrane (PubMed:[24904167](#), PubMed:[25315773](#), PubMed:[26567335](#), PubMed:[32493773](#)). Major PS-flippase in immune cell subsets. In erythrocyte plasma membrane, it is required to maintain PS in the inner leaflet preventing its exposure on the surface. This asymmetric distribution is critical for the survival of erythrocytes in circulation since externalized PS is a phagocytic signal for erythrocyte clearance by splenic macrophages (PubMed:[26944472](#)). Required for B cell differentiation past the pro-B cell stage (By similarity). Seems to mediate PS flipping in pro-B cells (By similarity). May be involved in the transport of cholestatic bile acids (By similarity).

Cellular Location

Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Early endosome membrane; Multi-pass membrane protein. Recycling endosome membrane; Multi-pass membrane protein. Note=Efficient exit from the endoplasmic reticulum requires the presence of TMEM30A. Internalized via clathrin-dependent endocytosis in response to Ca^{2+} signaling induced by G-protein coupled serotonin and histamine receptors

Tissue Location


Widely expressed.

ATP11C Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


ATP11C Antibody (Center) - Images




ATP11C Antibody (Center) (Cat.#AP5446c) western blot analysis in mouse lung tissue lysates (35ug/lane).This demonstrates the ATP11C antibody detected ATP11C protein (arrow).



ATP11C Antibody (Center) (Cat.#AP5446c) western blot analysis in CHO cell line lysates (35ug/lane).This demonstrates the ATP11C antibody detected the ATP11C protein (arrow).



ATP11C Antibody (Center) (Cat. #AP5446c) immunohistochemistry analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ATP11C Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



ATP11C Antibody (Center) (Cat. #AP5446c) flow cytometric analysis of 293 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

ATP11C Antibody (Center) - Background

The function of ATP11C remains unknown.

ATP11C Antibody (Center) - References

Matsuoka, S., et al. Science 316(5828):1160-1166(2007)
Andrew Nesbit, M., et al. Genomics 84(6):1060-1070(2004)
Halleck, M.S., et al. Physiol. Genomics 1(3):139-150(1999)
Yokoi, H., et al. Genomics 20(3):404-411(1994)

ATP11C Antibody (Center) - Citations

- [ATP11C targets basolateral bile salt transporter proteins in mouse central hepatocytes.](#)