

**ATP11C Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP5446c**

**Specification**

---

**ATP11C Antibody (Center) - Product Information**

Application	FC, IHC-P, WB,E
Primary Accession	<a href="#">Q8NB49</a>
Other Accession	<a href="#">NP_001010986.1</a>
Reactivity	Human, Hamster, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	129477
Antigen Region	589-616

**ATP11C Antibody (Center) - Additional Information**

**Gene ID** 286410

**Other Names**

Phospholipid-transporting ATPase IG, ATPase IQ, ATPase class VI type 11C, P4-ATPase flippase complex alpha subunit ATP11C, ATP11C, ATPIG, ATPIQ

**Target/Specificity**

This ATP11C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 589-616 amino acids from the Central region of human ATP11C.

**Dilution**

FC~~1:10~50

IHC-P~~1:50~100

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ATP11C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**ATP11C Antibody (Center) - Protein Information**

**Name** ATP11C {ECO:0000303|PubMed:26944472}

**Synonyms** ATPIG, ATPIQ

**Function** Catalytic component of a P4-ATPase flippase complex which catalyzes the hydrolysis of ATP coupled to the transport of aminophospholipids, phosphatidylserines (PS) and phosphatidylethanolamines (PE), from the outer to the inner leaflet of the plasma membrane (PubMed:[24904167](#), PubMed:[25315773](#), PubMed:[26567335](#), PubMed:[32493773](#)). Major PS-flippase in immune cell subsets. In erythrocyte plasma membrane, it is required to maintain PS in the inner leaflet preventing its exposure on the surface. This asymmetric distribution is critical for the survival of erythrocytes in circulation since externalized PS is a phagocytic signal for erythrocyte clearance by splenic macrophages (PubMed:[26944472](#)). Required for B cell differentiation past the pro-B cell stage (By similarity). Seems to mediate PS flipping in pro-B cells (By similarity). May be involved in the transport of cholestatic bile acids (By similarity).

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein. Endoplasmic reticulum membrane; Multi-pass membrane protein. Early endosome membrane; Multi-pass membrane protein. Recycling endosome membrane; Multi-pass membrane protein. Note=Efficient exit from the endoplasmic reticulum requires the presence of TMEM30A. Internalized via clathrin-dependent endocytosis in response to  $Ca^{2+}$  signaling induced by G-protein coupled serotonin and histamine receptors

#### **Tissue Location**


Widely expressed.

### **ATP11C Antibody (Center) - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


### **ATP11C Antibody (Center) - Images**




ATP11C Antibody (Center) (Cat.#AP5446c) western blot analysis in mouse lung tissue lysates (35ug/lane).This demonstrates the ATP11C antibody detected ATP11C protein (arrow).



ATP11C Antibody (Center) (Cat.#AP5446c) western blot analysis in CHO cell line lysates (35ug/lane).This demonstrates the ATP11C antibody detected the ATP11C protein (arrow).



ATP11C Antibody (Center) (Cat. #AP5446c) immunohistochemistry analysis in formalin fixed and paraffin embedded human prostate carcinoma followed by peroxidase conjugation of the secondary antibody and DAB staining. This data demonstrates the use of the ATP11C Antibody (Center) for immunohistochemistry. Clinical relevance has not been evaluated.



ATP11C Antibody (Center) (Cat. #AP5446c) flow cytometric analysis of 293 cells (bottom histogram) compared to a negative control cell (top histogram).FITC-conjugated goat-anti-rabbit secondary antibodies were used for the analysis.

#### ATP11C Antibody (Center) - Background

The function of ATP11C remains unknown.

#### **ATP11C Antibody (Center) - References**

Matsuoka, S., et al. Science 316(5828):1160-1166(2007)  
Andrew Nesbit, M., et al. Genomics 84(6):1060-1070(2004)  
Halleck, M.S., et al. Physiol. Genomics 1(3):139-150(1999)  
Yokoi, H., et al. Genomics 20(3):404-411(1994)

#### **ATP11C Antibody (Center) - Citations**

- [ATP11C targets basolateral bile salt transporter proteins in mouse central hepatocytes.](#)