

RNF39 Polyclonal Antibody
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP54485**Specification**

RNF39 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	O9H2S5
Reactivity	Rat, Dog
Host	Rabbit
Clonality	Polyclonal
Calculated MW	45525

RNF39 Polyclonal Antibody - Additional Information**Gene ID** 80352**Other Names**

RING finger protein 39, Protein HZFW, RNF39, HZFW

Format

0.01M TBS(pH7.4), 0.09% (W/V) sodium azide and 50% Glyce

Storage

Store at -20 °C for one year. Avoid repeated freeze/thaw cycles. When reconstituted in sterile pH 7.4 0.01M PBS or diluent of antibody the antibody is stable for at least two weeks at 2-4 °C.

RNF39 Polyclonal Antibody - Protein Information**Name** RNF39**Synonyms** HZFW**Function**

May play a role in prolonged long term-potential (LTP) maintenance.

Cellular Location

Cytoplasm.

Tissue Location


Expressed in testis..

RNF39 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RNF39 Polyclonal Antibody - Images




Protein: Raji lysate at 40ug;

Primary: rabbit Anti-RNF39 (bs-11393R) at 1:300;

Secondary: HRP conjugated Goat-Anti-rabbit IgG(bs-0295G-HRP) at 1: 5000;

Predicted band size: 38/45 kD

Observed band size: 38 kD



Paraformaldehyde-fixed, paraffin embedded (mouse brain tissue); Antigen retrieval by boiling in sodium citrate buffer (pH6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37°C for 30min; Antibody incubation with (RNF39) Polyclonal Antibody, Unconjugated (bs-11393R) at 1:400 overnight at 4°C, followed by operating according to SP Kit(Rabbit) (sp-0023) instructions and DAB staining.